

DHI 3 Algae and Sediment Model

MIKE ECO Lab Template

Scientific Description



DHI headquarters

Agern Allé 5
DK-2970 Hørsholm
Denmark

+45 4516 9200 Telephone

mike@dhigroup.com

www.mikepoweredbydhi.com

Company Registration No.: DK36466871

CONTENTS

DHI 3 Algae and Sediment Model MIKE ECO Lab Template Scientific Description

| | | |
|----------|--|----------|
| 1 | Introduction | 1 |
| 2 | Applications | 3 |
| 3 | Mathematical Formulations | 5 |
| 3.1 | Vertical light penetration | 8 |
| 3.2 | Production of autotrophs | 9 |
| 3.3 | Differential equations pelagic state variables | 12 |
| 3.3.1 | PC1: Flagellate C, g C m ⁻³ | 12 |
| 3.3.2 | PC2: Diatom C, g C m ⁻³ | 15 |
| 3.3.3 | PC3: Cyanobacteria C, g C m ⁻³ | 17 |
| 3.3.4 | PN1: Flagellate N, g N m ⁻³ | 20 |
| 3.3.5 | PN2, Diatom N, g N m ⁻³ | 22 |
| 3.3.6 | PN3, Cyanobacteria N, g N m ⁻³ | 23 |
| 3.3.7 | PP1, Flagellate P, g P m ⁻³ | 25 |
| 3.3.8 | PP2, Diatom P, g P m ⁻³ | 26 |
| 3.3.9 | PP3, Cyanobacteria P, g P m ⁻³ | 28 |
| 3.3.10 | PSi2, Diatom Si, g Si m ⁻³ | 29 |
| 3.3.11 | CH, Chlorophyll, g m ⁻³ | 31 |
| 3.3.12 | ZC, zooplankton, g C m ⁻³ | 33 |
| 3.3.13 | DC, Detritus C, g C m ⁻³ | 34 |
| 3.3.14 | DN, Detritus N, g N m ⁻³ | 36 |
| 3.3.15 | DP, Detritus P, g P m ⁻³ | 38 |
| 3.3.16 | DSi, Detritus Si, g Si m ⁻³ | 40 |
| 3.3.17 | NH4, Total ammonia, g N m ⁻³ | 41 |
| 3.3.18 | NO3, Nitrate, g N m ⁻³ | 45 |
| 3.3.19 | H2S, Hydrogen Sulphide, g S m ⁻³ | 47 |
| 3.3.20 | IP, Phosphate (PO4-P), g P m ⁻³ | 49 |
| 3.3.21 | IP, Phosphate (PO4-P), g P m ⁻³ | 52 |
| 3.3.22 | DO, Oxygen, g O2 m ⁻³ | 53 |
| 3.3.23 | CDOC, Coloured refractory DOC, g C m ⁻³ | 57 |
| 3.3.24 | CDON, Coloured refractory DON, g N m ⁻³ | 58 |
| 3.3.25 | CDOP, Coloured refractory DOP, g P m ⁻³ | 59 |
| 3.3.26 | LDOC, Labile DOC, g C m ⁻³ | 60 |
| 3.3.27 | LDON, Labile DON, g N m ⁻³ | 62 |
| 3.3.28 | LDOP, Labile DOP, g P m ⁻³ | 63 |
| 3.4 | Differential Equation Sediment State Variables..... | 65 |
| 3.4.1 | SSi, Sediment, bio-available Silicate, g Si m ²³ | 65 |
| 3.4.2 | KDOX, depth of NO3 penetration in sediment, m..... | 66 |
| 3.4.3 | KDO2, DO penetration in sediment, m..... | 67 |
| 3.4.4 | SOC, Sediment organic C, g C m ⁻² | 68 |
| 3.4.5 | SON, Bio-available organic N in sediment, g N m ⁻² | 70 |
| 3.4.6 | SOP, Bio-available organic P in sediment, g P m ⁻² | 72 |
| 3.4.7 | FESP, PO4 adsorbed to oxidised iron in sediment, g P m ⁻² | 74 |

| | | |
|----------|---|------------|
| 3.4.8 | SNH, Sediment pore water NH ₄ , g N m ⁻² | 75 |
| 3.4.9 | SNO ₃ , NO ₃ in sediment pore water, layer (0 - k _{do2}), g N m ⁻² | 76 |
| 3.4.10 | SIP, PO ₄ in sediment pore water, g P m ⁻² | 79 |
| 3.4.11 | SH ₂ S, Reduced substances in sediment, g S m ⁻² | 80 |
| 3.4.12 | SPI _M , Immobilised sediment P, g P m ⁻² | 82 |
| 3.4.13 | SN _{IM} , Immobilised sediment N by denitrification & burial, g N m ⁻² | 83 |
| 3.4.14 | SN _{IM} , Immobilised sediment N by denitrification & burial, g N m ⁻² | 84 |
| 3.5 | Help Processes | 85 |
| 3.5.1 | The P1 processes listed in alphabetic order | 85 |
| 3.5.2 | Auxiliary (A) processes listed in alphabetic order | 91 |
| | | |
| 4 | Data Requirements..... | 131 |
| | | |
| 5 | References | 133 |

1 Introduction

MIKE ECO Lab is a numerical lab for Ecological Modelling. It is a generic and open tool for customising aquatic ecosystem models to describe for instance water quality and eutrophication. DHI's expertise and know how concerning ecological modelling has been collected in predefined ecosystem descriptions (MIKE ECO Lab templates) to be loaded and used in MIKE ECO Lab. So the MIKE ECO Lab templates describe physical, chemical and biological processes related to environmental problems and water pollution. The following is a description of the DHI 3 algae and sediment model.

The DHI 3 algae and sediment template is used in investigations of eutrophication effects where different algae species and sediment pools of nutrients are essential and as an instrument in environmental impact assessments for such ecosystems. The 3 algae and sediment modelling can be applied in environmental impact assessments considering:

- Pollution sources such as domestic and industrial sewage and agricultural run-off
- Cooling water outlets from power plants resulting in excess temperatures
- Physical conditions such as sediment loads and change in bed topography affecting especially the benthic vegetation
- Evaluation of action plans related to nutrient reductions
- Risk evaluation in connection to potential harmful algae blooms

The aim of using 3 algae and sediment modelling as an instrument in environmental impact assessment studies is to obtain, most efficiently in relation to economy and technology, the optimal solution with regards to ecology and the human environment.

The 3 algae and sediment model describes nutrient cycling including internal loadings from sediment pools of nutrient, phytoplankton and zooplankton growth, in addition to simulating oxygen conditions.

The model results describe the concentrations of phytoplankton, chlorophyll-a, zooplankton, organic matter (detritus), organic and inorganic nutrients, oxygen and the area-based sediment pools of nitrogen and phosphorous over time. In addition to this, a number of derived variables are stored: primary production, total nitrogen and phosphorus concentrations, sediment oxygen demand and Secchi disc depth.

The 3 algae and sediment template is integrated with the advection-dispersion module, which describes the physical transport processes at each grid-point covering the area of interest. Other data required are concentrations at model boundaries, flow and concentrations from pollution sources, water temperature and influx of light, etc.

2 Applications

The eutrophication template can be applied in a range of environmental investigations:

- Studies where the effects of alternative nutrient loading situations are compared and/or different waste water treatment strategies are evaluated.
- Studies of oxygen depletion.
- Studies of the effects of the discharge of cooling water.
- Comparisons of the environmental consequences of different construction concepts for harbours, bridges, etc.
- Evaluation of the environmental consequences of developing new urban and industrial areas.
- Evaluation of action plans related to nutrient reductions and long term effects of reduction scenarios.
- Risk evaluation in connection to potential harmful algae blooms.

3 Mathematical Formulations

The MIKE 21/3 ECO Lab is coupled to the MIKE 21/3 AD module in order to simulate the simultaneous processes of transport, dispersion and biological/biochemical processes.

The 3 algae and sediment model includes state variables for 3 pelagic algae groups, nutrients, oxygen, hydrogen sulphide and sediment pools of C, N and P as well as a number of sediment state variables..

Table 3.1 Pelagic state variables

| Name | Comment | Unit |
|------|--|----------------------------------|
| PC1 | Flagellate C | g C m ⁻³ |
| PC2 | Diatom C | g C m ⁻³ |
| PC3 | Cyanobacteria C | g C m ⁻³ |
| PN1 | Flagellate N | g N m ⁻³ |
| PN2 | Diatom N | g N m ⁻³ |
| PN3 | Cyanobacteria N | g N m ⁻³ |
| PP1 | Flagellate P | g P m ⁻³ |
| PP2 | Diatom P | g P m ⁻³ |
| PP 3 | Cyanobacteria P | g P m ⁻³ |
| Psi | Diatom Si | g Si m ⁻³ |
| CH | Chlorophyll-a | g Chl m ⁻³ |
| ZC | Zooplankton C | g C m ⁻³ |
| DC | Detritus C | g C m ⁻³ |
| DN | Detritus N | g N m ⁻³ |
| DP | Detritus P | g P m ⁻³ |
| DSi | Detritus Si | g Si m ⁻³ |
| NH4 | Total ammonia (NH ₄) | g N m ⁻³ |
| NO3 | Nitrate+ nitrite | g N m ⁻³ |
| H2S | Hydrogen Sulphide (H ₂ S) | g S m ⁻³ |
| IP | Inorganic Phosphorous (PO ₄) | g P m ⁻³ |
| Si | Silicate Si | g Si m ⁻³ |
| DO | Dissolved Oxygen | g O ₂ m ⁻³ |
| CDOC | Coloured refractory DOC | g C m ⁻³ |
| CDON | Coloured refractory DON | g N m ⁻³ |
| CDOP | Coloured refractory DOP | g P m ⁻³ |
| LDOC | Labile DOC | g C m ⁻³ |
| LDON | Labile DON | g N m ⁻³ |
| LDOP | Labile DOP | g P m ⁻³ |

Table 3.2 Sediment state variables

| Name | Comment | Unit |
|------|--|----------|
| SSi | Sediment biological available Silicate | g Si m-2 |
| KDOX | Oxidised layer, depth of NO3 penetration in sediment | m |
| KDO2 | DO penetration into sediment | m |
| SOC | Sediment organic C | g C m-2 |
| SON | Sediment organic N | g N m-2 |
| SOP | Sediment organic P | g P m-2 |
| FESP | Sediment iron adsorbed PO4 | g P m-2 |
| SNH | Sediment pore water NH4 | g N m-2 |
| SNO3 | NO3-N in Surface sediment pore water, layer (0 - kdo2) | g N m-2 |
| SIP | Sediment pore water PO4 | g P m-2 |
| SH2S | Sediment reduced substances as (H2S) | g S m-2 |
| SPIM | Immobilised P in sediment | g P m-2 |
| SNIM | Sediment immobilised N by denitrification & burial | g N m-2 |
| SCIM | Sediment immobilised C by mineralisation & burial of SOC | g C m-2 |

Table 3.3 Additional State variables for mass considerations

| Name | Comment | Unit |
|------------|--|----------|
| sum_PRPC | Sum of PC production | g C m-2 |
| sum_CminW | Sum of pelagic C mineralisation | g C m-2 |
| sum_minSOC | Sum of SOC mineralisation | g C m-2 |
| sum_DEPON | Sum of atmospheric deposition of N | g N m-2 |
| Sum_Nfix | Sum of cyanobacterial N fixation | g N m-2 |
| sum_DENW | Sum of denitrification in water column | g N m-2 |
| sum_Nflux | Sum of N flux sediment- water | g N m-2 |
| sum_rdenit | Sum of sediment denitrification | g N m-2 |
| sum_DEPOP | Sum of atmospheric deposition of P | g P m-2 |
| sum_Pflux | Sum of P flux sediment-water | g P m-2 |
| sum_rear | Sum of reaeration | g O2 m-2 |
| sum_ODSC | Sum of sediment O2 respiration | g O2 m-2 |
| sum_RSH2S | Sum of H2S production in sediment | g S m-2 |

The first 28 components or state variables (pelagic system) are moveable and treated in both the MIKE 21/3 AD and the MIKE 21/3 ECO Lab module. The additional components have a fixed nature belonging to the benthic system.

The processes and transfer of carbon, nitrogen and phosphorus in the Eutrophication model system is illustrated in Figure 3.1. Also included in the model is an oxygen balance.

The processes describing the variations of the components in time and space are dependent on external factors such as the salinity, water temperature, the light influx, and the discharges.

The salinity and water temperature can be results of MIKE 21/3 AD simulations or be user specified values. The first possibility is especially relevant for cooling water investigations whereas the latter possibility often is used in areas where only natural variations in temperature are seen.

The mathematical formulations of the biological and chemical processes and transformations for each state variable are described one by one below. The differential equations are 1st order, ordinary and coupled.




Figure 3.1 The simplified flow diagram of the fluxes of carbon, nitrogen and phosphorus in the eutrophication model.

3.1 Vertical light penetration

Light is essential for growth of all plants, including the pelagic. The vertical light penetration can be described by an exponential decay with depth which is dependent on a light extinction K_d , which either can be described as with light extinction constants (k_{dx}) multiplied by concentrations of light extinction concentration (Chlorophyll (CH, $g\ m^{-3}$)), detritus (DC, $g\ C\ m^{-3}$), dissolved organic matter (CDOC, $g\ C\ m^{-3}$), inorganic matter (SS, $g\ m^{-3}$) and water (k_{bla} , m^{-1}) or it can be described as a function of scattering (b , m^{-1}) and absorption (a , m^{-1}) of light.

Vertical light penetration with depth (z , m) in the water column:

$$I_z = I_0 * e^{-K_{dx} * z} \text{ mol photons } m^2\ d^{-1} \quad (3.1)$$

Where K_{dx} can be either K_{d1} or K_{d2} :

$$K_{d1} = K_{chl} * CH + K_{dc} * DC + k_{cdoc} * CDOC + k_{ss} * SS + k_{bla}, m^{-1} \quad (3.2)$$

Or:

$$K_{d2} = \sqrt{a^2 + 0.256 * a * b}, m^{-1} \quad (3.3)$$

The absorption of light is mainly associated to particulate and dissolved organic matter whereas the scattering is mainly associated to particulate inorganic matter.

Light absorption, where the notation K_{xa} stand for light absorption constant of component x:

$$a = K_{chl_a} * CH + K_{dc_a} * DC + K_{cdoc_a} * CDOC + K_{ss_a} * SS + k_{bla}, m^{-1} \quad (3.4)$$

Light scattering from phytoplankton and fine suspended inorganic matter can be describes as power functions of CH and SS:

$$b = bkch * CH^{ekch} + bkss * SS^{ekss}, m^{-1} \quad (3.5)$$

Where the light scattering constants ($bkch$, $bkss$ in m^2g^{-1}) and exponents ($ekch$, $ekss$) are for chlorophyll and inorganic suspended matter, respectively.

The present ecological model do not simulate resuspension of (fine) sediment, therefore SS is not dynamically simulated. Resuspension is most pronounced on shallow waters below 5-10 m. The user should therefore consider the need for either including measured SS concentrations or modelled concentrations of SS by a sediment transport model (MIKE by DHI 2011a). On shallow waters (like lagoons) the EU-MT MIKE ECO Lab template can be used. This template includes resuspension of and transport of fine sediment and combine it with a description of nutrients (N, P) one phytoplankton group, one macroalgae, one rooted macrophyte (eelgrass) and microbenthic algae (Rasmussen E. K. et al. 2009).

The present model however calculates dynamically the concentration of chlorophyll (CH), detritus carbon (DC) and refractory or coloured dissolved organic C (CDOC). The missing resuspension of SS is minimal if used on set up with waters above 10 m depth, like the Baltic Sea.

3.2 Production of autotrophs

The template includes 3 pelagic autotrophs (flagellates, diatoms and cyanobacteria). The production is based on daily dose of photosynthetic active light (PAR, mol photons m²d⁻¹) light resulting in a net production.

The differential equation includes a net production, sedimentation, buoyancy (flagellate & cyanobacteria) and mortality by grazing and nutrient limitation (nutrient stress).

$$\frac{dx}{dt} = \text{gross production} - \text{mortality} - \text{sedimentation} + \text{boyancy} \quad (3.6)$$

The net production is determined by light ($flight(i)$), temperature ($ftemp(T)$) and nutrient availability ($fnut(N,P, (Si \text{ diatoms}))$). μ_T is the temperature corrected max specific growth (d⁻¹) and X is the biomass (g C m⁻³ or g C m⁻²)

$$\text{net production} = \mu_T * flight(i) * ftemp(T) * fnut(N, P, (Si)) * FAC * RD * X \quad (3.7)$$

Where:

| Name | Comment | Unit |
|---------|--|--|
| μ | Max specific net growth rate (12 h light/12 h dark) at 20 °C | d ⁻¹ |
| i | Light (PAR) dose | mol photon m ⁻² d ⁻¹ |
| T | Temperature | °C |
| N,P, Si | Internal concentrations of N, P and SI in algae | g nutrient g C ⁻¹ |
| FAC | Correction of dark reaction (growth) | n.u. |
| RD | Relative day length, function of latitude, 1 at 12 h light | n.u |

Temperature is an important direct or indirect regulator of many processes. Two types of temperature functions are used, Arrhenius or Lassiter functions.

The Arrhenius function increases the process exponentially with temperature; whereas the Lassiter function have an optimum temperature from which the process decline towards zero.

In the present template Arrhenius relations are used to describe the max specific growth rates, as the template is used in waters where the temperature rarely exceeds 20 °C. Further at increasing temperatures the plankton community will have a tendency to adapt to the higher temperature by change of species composition.

The user is encouraged to consider the feasibility to change from Arrhenius to a Lassiter temperature regulation of the max specific growth rates if needed. Both Arrhenius and Lassiter expressions are bullied in function in MIKE ECO Lab see (MIKE by DHI 2011b).

Lassiter functions are used to temperature regulate the max specific growth rates. In contrast to the Arrhenius function The Lassiter function include an optimum temperature above which the function will decline. Arrhenius functions are used to regulate the specific growth of phytoplankton or macrophytes in areas normally having summer temperatures well above 20 °C.

It is still recommended to use Arrhenius functions for temperature regulation of respiration processes; however the user might consider to increase the reference temperature for 20 °C, if data or references justify this.

Lassiter:

$$L(T) = p_{opt} * e^{K2*(T-T_{opt})} * \left(\frac{(T_{max}-T)}{(T_{max}-T_{opt})} \right)^{K2*(T_{max}-T_{opt})}, d^{-1} \quad (3.8)$$

Arrhenius 20 °C:

$$A(T) = \theta^{(T-20)}, d^{-1} \quad (3.9)$$

Where:

- T: Temperature °C
- popt: max growth at T_{opt}, d-1
- Topt: Optimum temperature °C
- Tmax: Maximum temperature °C
- Θ: Theta constant Arrhenius function
- K2: constant




Figure 3.2 Arrhenius at 5 and 20 °C (Θ 1.04) and Lassiter function at T_{opt} at 12 and 18 °C (T_{max} 30 °C, K2 0.4)

The nutrient regulates the growth of all autotrophs. Two different nutrient regulators of the growth are used. A Droop kinetic (Droop 1973, Droop 1975) is used for autotrophs having internal nutrient pools (flagellates, diatoms, cyanobacteria). A Monod kinetics (Monod J. 1949) is used to describe the uptake of inorganic N, P and Si from the water into plankton.

Further cyanobacteria has the ability to N fixation in situation where the internal N:C ratio is low and the internal P:C is above average.

Nutrient regulation of primary production of phytoplankton (flagellates and cyanobacteria):

$$fnut(N, P)_{pm} = \frac{2}{\frac{1}{f(N)_{pm}} + \frac{1}{f(P)_{pm}}} \tag{3.10}$$

In the expression for diatoms Si is included:

$$fnut(N, P, Si)_{pm} = \frac{3}{\frac{1}{f(N)_{pm}} + \frac{1}{f(P)_{pm}} + \frac{1}{f(Si)_{pm}}} \tag{3.11}$$

Droop kinetics used for N modified after (Nyholm1978, Nyholm 1979) is used to regulate growth of phytoplankton:

$$f(N)_{pm} = \frac{\frac{PN}{PC} - PNmin}{PNmax - PNmin} \tag{3.12}$$

The same formulation is used for diatoms.

Droop kinetics used for P modified after (Nyholm1978, Nyholm 1979) is used to regulate growth of phytoplankton:

$$f(P)_{pm} = \frac{\left(\frac{PP}{PC} - PPmin\right) * (Kc + PPmax - PPmin)}{(PPmax - PPmin) * \left(Kc + \frac{PP}{PC} - PPmin\right)} \tag{3.13}$$

Where:

| Name | Comment | Unit |
|-------|-------------------------------------|-----------|
| PC | Phytoplankton C | g C m-3 |
| PN | Phytoplankton N | g N m-3 |
| PP | Phytoplankton P | g P m-3 |
| PNmin | Minimum N:C ratio for phytoplankton | g N g C-1 |
| PNmax | Maximum N:C ratio for phytoplankton | g N g C-1 |
| PPmin | Minimum P:C ratio for phytoplankton | g P g C-1 |
| PPmax | Maximum P:C ratio for phytoplankton | g P g C-1 |

3.3 Differential equations pelagic state variables

3.3.1 PC1: Flagellate C, g C m⁻³

$$dPC1/dt = PRPC1 - GRPC1 - DEPC1 - SEPC1 - BUOYC1 \quad (3.14)$$

Where:

| Process | Comment | Unit |
|---------|----------------------------------|-------------------------------------|
| PRPC1 | Net production flagellate carbon | g C m ⁻³ d ⁻¹ |
| GRPC1 | Grazing of flagellate carbon | g C m ⁻³ d ⁻¹ |
| DEPC1 | Death of flagellate carbon | g C m ⁻³ d ⁻¹ |
| SEPC1 | Settling of flagellate carbon | g C m ⁻³ d ⁻¹ |
| BUOYC | Flagellate upward movement | g C m ⁻³ d ⁻¹ |

PRPC1: Net Production flagellate carbon, g C m⁻³ d⁻¹

$$PRPC1 = mntp1 * myfi1 * fac * rd * PC1 \quad (3.15)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-----------------|--------|
| mntp1 | N, P & temperature corrected max. net growth rate | d ⁻¹ | A |
| myfi1 | Light function Flagellate, | n.u. | A |
| fac | Phytoplankton, Correction for dark reaction | n.u. | C |
| rd | Relative daylength, f(latitude, day,month,year) | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

GRPC1: Grazing of phytoplankton (Flagellate) carbon, g C m⁻³d⁻¹

$$GRPC1 = \frac{kedib1 * MAX(PC1 - 0.001)}{MAX(0.001, kedib1 * PC1 + kedib2 * PC2 + kedib3 * PC3)} * mgpc * ZC \tag{3.16}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|---------------------|--------|
| Kedib1 | Edible fraction of Flagellate | n.u. | C |
| Kedib2 | Edible fraction of Diatoms | n.u. | C |
| Kedib3 | Edible fraction of Cyanobacteria | n.u. | C |
| PC1 | Flagellate C | g C m ⁻³ | S |
| PC2 | Diatom C | g C m ⁻³ | S |
| PC3 | Cyanobacterie C | g C m ⁻³ | S |
| mgpc | Temperature & food corrected grazing rate | d ⁻¹ | A |
| ZC | Zooplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

DEPC1: Death of phytoplankton (flagellate) carbon, g C m⁻³d⁻¹

$$DEPC1 = kdma * mnl1 * PC1 \tag{3.17}$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|-----------------|--------|
| kdma | Specific death rate phytoplankton | d ⁻¹ | C |
| mnl1 | Nutrient dependent death factor, flagellate | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

SEPC1: Settling of phytoplankton (flagellate) carbon, g C m⁻³d⁻¹

Phytoflagellates has the ability of vertical movement in the water column optimising their ability to pick up nutrient and gain light. During nutrient limitation the flagellates is assumed to seek down to the pycnocline to pick up nutrient, and in case they are not nutrient limited they are assumed to stay in the photic zone.

In the present model nutrient limitation, in term of a low PN/PC and or PP/PC ratio, enhance the sedimentation rate. PN/PC and PP/PC ratios close to maximum N and P content in the algae result in a reduction of the sedimentation rate. The nutrient regulation of the sedimentation rate is expressed in the auxiliary *sed1*.

Light is also regulating the sedimentation rate. At high light dozes the sedimentation is accelerated at medium light dozes sedimentation is *m_{spc1}* and at low light dozes the sedimentation decreases. This light regulation is expressed in the auxiliary *fiz*.

$$SEPC1 = \frac{m_{spc1}}{dz} * sed1 * fiz * PC1 \quad (3.18)$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|---|-------------------|--------|
| m _{spc1} | Sedimentation rate flagellate phytoplankton | m d ⁻¹ | A |
| Dz | Height of actual water layer | m | F |
| sed1 | N & P regulation of sedimentation. flagellate | n.u. | A |
| Fiz | Light factor for PC1 & PC3 sedimentation | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

BUOYC1: Flagellate upward movement, g C m⁻³d⁻¹

The vertical upward movement by the phytoflagellates is described as a function of light doze and the algae's nutrient condition expressed in the auxiliary *buoy1*. An upward vertical movement is enhanced by a good nutrient condition and a low light doze.

$$BUOYC1 = \frac{m_{spc1} * buoy1 * PC1}{dz} \quad (3.19)$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|--|-------------------|--------|
| m _{spc1} | Sedimentation rate flagellate phytoplankton | m d ⁻¹ | A |
| buoy1 | N & P & light upward movement function, flagellate | n.u. | A |
| Dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.2 PC2: Diatom C, g C m⁻³

$$\frac{dPC2}{dt} = PRPC2 + GRPC2 - DEPC2 - SEPC2 \tag{3.20}$$

Where:

| Process | Comment | Unit |
|---------|------------------------------|-------------------------------------|
| PRPC2 | Net production diatom carbon | g C m ⁻³ d ⁻¹ |
| GRPC2 | Grazing of diatom carbon | g C m ⁻³ d ⁻¹ |
| DEPC2 | Death of diatom carbon | g C m ⁻³ d ⁻¹ |
| SEPC2 | Settling of diatom carbon | g C m ⁻³ d ⁻¹ |

PRPC2: Net Production phytoplankton carbon, g C m⁻³d⁻¹

$$PRPC2 = mntp2 * myfi * fac * rd * PC2 \tag{3.21}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-----------------|--------|
| mntp2 | N, P, Si & temperature corrected max. net growth rate | d ⁻¹ | A |
| myfi2 | Light function Diatom | n.u. | A |
| fac | Phytoplankton, correction for dark reaction | n.u. | C |
| rd | Relative daylength, f(latitude, day,month,year) | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

GRPC2: Grazing of phytoplankton (Diatom) carbon, g C m⁻³d⁻¹

$$GRPC2 = \frac{kedib2 * MAX(PC2 - 0.001)}{MAX(0.001, kedib1 * PC1 + kedib2 * PC2 + kedib3 * PC3) * ZC} * mgpc \quad (3.22)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|---------------------|--------|
| Kedib1 | Edible fraction of Flagellate | n.u. | C |
| Kedib2 | Edible fraction of Diatoms | n.u. | C |
| Kedib3 | Edible fraction of Cyanobacteria | n.u. | C |
| PC1 | Flagellate C | g C m ⁻³ | S |
| PC2 | Diatom C | g C m ⁻³ | S |
| PC3 | Cyanobacterie C | g C m ⁻³ | S |
| mgpc | Temperature & food corrected grazing rate | d ⁻¹ | A |
| ZC | Zooplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

DEPC2: Death of diatom carbon, g C m⁻³d⁻¹

$$DEPC2 = kdma * mnl2 * PC2 \quad (3.23)$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|-----------------|--------|
| kdma | Specific death rate phytoplankton | d ⁻¹ | C |
| mnl2 | Nutrient dependent death factor, diatom | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

SEPC2: Settling of diatom carbon, g C m⁻³d⁻¹

$$SEPC2 = mspc2 * \frac{mnl2}{dz} * PC2 \tag{3.24}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|--|-------------------|--------|
| mspc2 | Sedimentation rate diatom phytoplankton | m d ⁻¹ | A |
| mnl2 | Nutrient function, sedimentation & death, diatom | n.u. | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.3 PC3: Cyanobacteria C, g C m⁻³

$$dPC3/dt = PRPC3 - GRPC3 - DEPC3 - SEPC3 - BUOYC3 \tag{3.25}$$

Where:

| Process | Comment | Unit |
|---------|-------------------------------------|-------------------------------------|
| PRPC3 | Net production cyanobacteria carbon | g C m ⁻³ d ⁻¹ |
| GRPC3 | Grazing of cyanobacteria carbon | g C m ⁻³ d ⁻¹ |
| DEPC3 | Death of cyanobacteria carbon | g C m ⁻³ d ⁻¹ |
| SEPC3 | Settling of cyanobacteria carbon | g C m ⁻³ d ⁻¹ |
| BUOYC3 | Cyanobacteria upward movement | g C m ⁻³ d ⁻¹ |

PRPC3: Net Production cyanobacteria carbon, g C m⁻³d⁻¹

$$PRPC3 = mntp3 * myfi3 * fp3sal * fac * rd * PC3 \quad (3.26)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|-----------------|--------|
| mntp3 | N, P & temperature corrected max. net growth rate | d ⁻¹ | A |
| myfi3 | Light function Cyanobacteria, | n.u. | A |
| fp3sal | Function for cyanobacteria dependency of salinity | n.u. | A |
| fac | Cyanobacteria, Correction for dark reaction | n.u. | C |
| rd | Relative daylength, f(latitude, day,month,year) | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

GRPC3: Grazing of cyanobacteria (Flagellate) carbon, g C m⁻³d⁻¹

$$GRPC3 = \frac{kedib3 * MAX(PC3 - 0.001)}{MAX(0.001, kedib1 * PC1 + kedib2 * PC2 + kedib3 * PC3) * ZC} * mgpc \quad (3.27)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|---------------------|--------|
| Kedib1 | Edible fraction of Cyanobacteria | n.u. | C |
| Kedib2 | Edible fraction of Diatoms | n.u. | C |
| Kedib3 | Edible fraction of Cyanobacteria | n.u. | C |
| PC1 | Flagellate C | g C m ⁻³ | S |
| PC2 | Diatom C | g C m ⁻³ | S |
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| mgpc | Temperature & food corrected grazing rate | d ⁻¹ | A |
| ZC | Zooplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

DEPC3: Death of cyanobacteria carbon, g C m⁻³d⁻¹

$$DEPC3 = kdma * \frac{mnl3}{fp3sal} * PC3 \tag{3.28}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|-----------------|--------|
| kdma | Specific death rate cyanobacteria | d ⁻¹ | C |
| mnl3 | Nutrient dependent death factor, cyanobacteria | n.u. | A |
| fp3sal | Function for cyanobacteria dependency of salinity | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

SEPC3: Settling of cyanobacteria carbon, g C m⁻³d⁻¹

Cyanobacteria has the ability of vertical movement in the water column optimising their ability to pick up nutrient and gain light. During nutrient limitation the cyanobacteria seek down to the pycnocline to pick up P nutrients, and in case they are not nutrient limited (P) they are assumed to stay in the photic zone.

In the present model nutrient limitation, in term of a low PN/PC and or PP/PC ratio, enhance the sedimentation rate. PN/PC and PP/PC ratios close to maximum N and P content in the algae result in a reduction of the sedimentation rate. The nutrient regulation of the sedimentation rate is expressed in the auxiliary **sed3**.

Light is also regulating the sedimentation rate. At high light dozes the sedimentation is accelerated at medium light dozes sedimentation is **m_{spc3}** and at low light dozes the sedimentation decreases. This light regulation is expressed in the auxiliary **fiz**.

$$SEPC3 = \frac{m_{spc3}}{dz} * sed3 * fiz * PC3 \tag{3.29}$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|--|-------------------|--------|
| m _{spc3} | Sedimentation rate cyanobacteria | m d ⁻¹ | A |
| Dz | Height of actual water layer | m | F |
| sed3 | N & P regulation of sedimentation. cyanobacteria | n.u. | A |
| fiz | Light factor for PC1 & PC3 sedimentation | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

BUOYC3: Cyanobacteria upward movement, g C m⁻³d⁻¹

The vertical upward movement by the cyanobacteria is described as a function of light doze and the algae's nutrient condition expressed in the auxiliary **buoy3**. An upward vertical movement is enhanced by a good nutrient condition and a low light doze.

$$BUOYC3 = \frac{mspc3 * buoy3 * PC3}{dz} \quad (3.30)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-----------------|--------|
| mspc3 | Sedimentation rate cyanobacteria | d ⁻¹ | A |
| Buoy3 | N & P & light upward movement function, cyanobacteria | n.u. | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.4 PN1: Flagellate N, g N m⁻³

$$\frac{dPN1}{dt} = UPNH1 + UPN31 - GRPN1 - DEPN1 - SEPN1 - BUOYN1 \quad (3.31)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| UPNH1 | Uptake of NH ₄ into flagellates N | g N m ⁻³ d ⁻¹ |
| UPN31 | Uptake of NO ₃ into flagellates N | g N m ⁻³ d ⁻¹ |
| GRPN1 | Grazing of flagellates N | g N m ⁻³ d ⁻¹ |
| DEPN1 | Death of flagellates N | g N m ⁻³ d ⁻¹ |
| SEPN1 | Settling of flagellates N | g N m ⁻³ d ⁻¹ |
| BUOYN1 | Upward movement flagellate N | g N m ⁻³ d ⁻¹ |

UPNH1: Uptake of NH₄ into flagellate N, g N m⁻³ d⁻¹

$$UPNH1 = MIN(unh1, pnma * PRPC1) \quad (3.32)$$

UPN31: Uptake of NO₃ into flagellate N, g N m⁻³ d⁻¹

$$UPN31 = MAX(0, MIN(un31, pnma * PRPC1 - UPNH1)) \quad (3.33)$$

GRPN1: Grazing of flagellate N, g N m⁻³ d⁻¹

$$GRPN1 = pn1pc1 * GRPC1 \quad (3.34)$$

DEPN1: Death of flagellate N, g N m⁻³ d⁻¹

$$DEPN1 = pn1pc1 * DEPC1 \quad (3.35)$$

SEPN1: Settling of flagellate N, g N m⁻³ d⁻¹

$$SEPN1 = pn1pc1 * SEPC1 \quad (3.36)$$

BUOYN1: Upward movement of PN1, g N m⁻³ d⁻¹

$$BUOYN1 = pn1pc1 * BUOYC1 \quad (3.37)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| pnma | Max. intracellular algae N | g N g C ⁻¹ | C |
| unh1 | potential NH ₄ uptake by flagellate | g N m ⁻³ d ⁻¹ | A |
| un31 | potential NO ₃ uptake by flagellate | g N m ⁻³ d ⁻¹ | A |
| pn1pc1 | Flagellate N:C ration | g N g C ⁻¹ | A |
| PC1 | Flagellate C | g C m ⁻³ | S |
| GRPC1 | Grazing of flagellate C | g C m ⁻³ d ⁻¹ | P |
| DEPC1 | Death of flagellate C | g C m ⁻³ d ⁻¹ | P |
| SEPC1 | Settling of flagellate C | g C m ⁻³ d ⁻¹ | P |
| BUOYC1 | Upward movement of flagellate C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.5 PN2, Diatom N, g N m⁻³

$$\frac{dPN2}{dt} = UPNH2 + UPN32 - GRPN2 - DEPN2 - SEPN2 \quad (3.38)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| UPNH2 | Uptake of NH ₄ into diatoms N | g N m ⁻³ d ⁻¹ |
| UPN32 | Uptake of NO ₃ into diatoms N | g N m ⁻³ d ⁻¹ |
| GRPN2 | Grazing of diatoms N | g N m ⁻³ d ⁻¹ |
| DEPN2 | Death of diatoms N | g N m ⁻³ d ⁻¹ |
| SEPN2 | Settling of diatoms N | g N m ⁻³ d ⁻¹ |

UPNH2: Uptake of NH₄ into diatom N, g N m⁻³ d⁻¹

$$UPNH2 = MIN(unh2, pnma * PRPC2) \quad (3.39)$$

UPN32: Uptake of NO₃ into diatom N, g N m⁻³ d⁻¹

$$UPN32 = MAX(0, MIN(un32, pnma * PRPC2 - UPNH2)) \quad (3.40)$$

GRPN2: Grazing of diatom N, g N m⁻³ d⁻¹

$$GRPN2 = pn2pc2 * GRPC2 \quad (3.41)$$

DEPN2: Death of diatom N, g N m⁻³ d⁻¹

$$DEPN2 = pn2pc2 * DEPC2 \quad (3.42)$$

SEPN2: Settling of diatom N, g N m⁻³ d⁻¹

$$SEPN2 = pn2pc2 * SEPC2 \quad (3.43)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| pnma | Max. intracellular algae N | g N g C ⁻¹ | C |
| unh2 | potential NH ₄ uptake by diatom | g N m ⁻³ d ⁻¹ | A |
| un32 | potential NO ₃ uptake by diatom | g N m ⁻³ d ⁻¹ | A |
| pn2pc2 | Diatom N:C ration | g N g C ⁻¹ | A |
| PC2 | Diatom C | g C m ⁻³ | S |
| GRPC2 | Grazing of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| SEPC2 | Settling of diatom C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.6 PN3, Cyanobacteria N, g N m⁻³

$$\frac{dPN3}{dt} = UPNH3 + UPN33 + NFIX - GRPN3 - DEPN3 - SEPN3 - BUOYN3 \quad (3.44)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| UPNH3 | Uptake of NH ₄ into cyanobacteria N | g N m ⁻³ d ⁻¹ |
| UPN33 | Uptake of NO ₃ into cyanobacteria N | g N m ⁻³ d ⁻¹ |
| NFIX | N fixation cyanobacteria | g N m ⁻³ d ⁻¹ |
| GRPN3 | Grazing of cyanobacteria N | g N m ⁻³ d ⁻¹ |
| DEPN3 | Death of cyanobacterias N | g N m ⁻³ d ⁻¹ |
| SEPN3 | Settling of cyanobacterias N | g N m ⁻³ d ⁻¹ |
| BUOYN3 | Upward movement cyanobacteria N | g N m ⁻³ d ⁻¹ |

UPNH3: Uptake of NH₄ into cyanobacteria N, g N m⁻³ d⁻¹

$$UPNH3 = MIN(unh3, pnma * PRPC3) \quad (3.45)$$

UPN33: Uptake of NO₃ into cyanobacteria N, g N m⁻³ d⁻¹

$$UPN33 = MAX(0, MIN(un33, pnma * PRPC3 - UPNH3)) \quad (3.46)$$

NFIX: N fixation by cyanobacteria N, g N m⁻³ d⁻¹

$$NFIX = IF (krednc - pn3pc3) < epsi THEN 0 ELSE knfix1 * tppc^{T-20} * \frac{MAX(0, krednc - pn3pc3)}{MAX(0, krednc - pn3pc3) + kqppn} * nfix1 * nfix2 \quad (3.47)$$

GRPN3: Grazing of cyanobacteria N, g N m⁻³ d⁻¹

$$GRPN3 = pn3pc3 * GRPC3 \quad (3.48)$$

DEPN3: Death of cyanobacteria N, g N m⁻³ d⁻¹

$$DEPN3 = pn3pc3 * DEPC3 \quad (3.49)$$

SEPN3: Settling of cyanobacteria N, g N m⁻³ d⁻¹

$$SEPN3 = pn3pc3 * SEPC3 \quad (3.50)$$

BUOYN3: Upward movement of PN3, g N m⁻³ d⁻¹

$$BUOYN3 = pn3pc3 * BUOYC3 \quad (3.51)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|---------------------------------------|--------|
| Pnma | Max. intracellular algae N | g N g C ⁻¹ | C |
| unh3 | potential NH ₄ uptake by cyanobacteria | g N m ⁻³ d ⁻¹ | A |
| un33 | potential NO ₃ uptake by cyanobacteria | g N m ⁻³ d ⁻¹ | A |
| pn3pc3 | Cyanobacteria N:C ration | g N g C ⁻¹ | A |
| krednc | Redfield ratio N:C | g N g C ⁻¹ | C |
| knfix1 | Max. N fixation, 20 °C, cyanobacteria | g N g C ⁻¹ d ⁻¹ | C |
| Tppc | Θ in Arrhenius temperature function | n.u. | C |
| T | Temperature | °C | F |
| Kqppn | Half saturation constant, N fixation | g N g C ⁻¹ | C |
| nfix1 | Function for N fixation (1 if PSU≤12 else 0) | n.u. | A |

| Name | Comment | Unit | Type*) |
|--------|---|-------------------------------------|--------|
| nfix2 | Function for N fixation (1 if $0 \leq \text{PSU} \leq 10$ else 0-1) | n.u. | A |
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| GRPC3 | Grazing of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| DEPC3 | Death of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| SEPC3 | Settling of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| BUOYC3 | Upward movement of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.7 PP1, Flagellate P, g P m⁻³

$$\frac{dPP1}{dt} = UPPP1 - GRPP1 - DEPP1 - SEPP1 - BUOYP1 \quad (3.52)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| UPPP1 | Uptake of PO ₄ into flagellates P | g P m ⁻³ d ⁻¹ |
| GRPP1 | Grazing of flagellates P | g P m ⁻³ d ⁻¹ |
| DEPP1 | Death of flagellates P | g P m ⁻³ d ⁻¹ |
| SEPP1 | Settling of flagellates P | g P m ⁻³ d ⁻¹ |
| BUOYP1 | Upward movement flagellate | g P m ⁻³ d ⁻¹ |

UPPP1: Uptake of PO₄ into flagellate P, g P m⁻³ d⁻¹

$$UPPP1 = \text{MIN}(upo1, ppma * PRPC1) \quad (3.53)$$

GRPP1: Grazing of flagellate P, g P m⁻³ d⁻¹

$$GRPP1 = pp1pc1 * GRPC1 \quad (3.54)$$

DEPP1: Death of flagellate P, g P m⁻³ d⁻¹

$$DEPP1 = pp1pc1 * DEPC1 \quad (3.55)$$

SEPP1: Settling of flagellate P, g P m⁻³ d⁻¹

$$SEPP1 = pp1pc1 * SEPC1 \quad (3.56)$$

BUOYP1: Upward movement of PP1, g P m⁻³ d⁻¹

$$BUOYP1 = pp1pc1 * BUOYC1 \quad (3.57)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| pnma | Max. intracellular algae P | g P g C ⁻¹ | C |
| upo1 | Potential PO ₄ uptake by flagellate | g P m ⁻³ d ⁻¹ | A |
| pp1pc1 | Flagellate P:C ration | g P g C ⁻¹ | A |
| PC1 | Flagellate C | g C m ⁻³ | S |
| GRPC1 | Grazing of flagellate C | g C m ⁻³ d ⁻¹ | P |
| DEPC1 | Death of flagellate C | g C m ⁻³ d ⁻¹ | P |
| SEPC1 | Settling of flagellate C | g C m ⁻³ d ⁻¹ | P |
| BUOYC1 | Upward movement of flagellate C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.8 PP2, Diatom P, g P m⁻³

$$\frac{dPP2}{dt} = UPPP2 - GRPP2 - DEPP2 - SEPP2 \quad (3.58)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| UPPP2 | Uptake of PO ₄ into diatoms P | g P m ⁻³ d ⁻¹ |
| GRPP2 | Grazing of diatoms P | g P m ⁻³ d ⁻¹ |
| DEPP2 | Death of diatoms P | g P m ⁻³ d ⁻¹ |
| SEPP2 | Settling of diatoms P | g P m ⁻³ d ⁻¹ |

UPPP2: Uptake of PO₄ into diatom P, g P g P m⁻³ d⁻¹

$$UPPP2 = \text{MIN}(upo2, ppma * PRPC2) \quad (3.59)$$

GRPP2: Grazing of diatom P, g P m⁻³ d⁻¹

$$GRPP2 = pp2pc2 * GRPC2 \quad (3.60)$$

DEPP2: Death of diatom P, g P m⁻³ d⁻¹

$$DEPP2 = pp2pc2 * DEPC2 \quad (3.61)$$

SEPP2: Settling of diatom P, g P m⁻³ d⁻¹

$$SEPP2 = pp2pc2 * SEPC2 \quad (3.62)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| pnma | Max. intracellular algae P | g P g C ⁻¹ | C |
| upo2 | Potential PO ₄ uptake by diatom | g P m ⁻³ d ⁻¹ | A |
| pp2pc2 | Diatom P:C ration | g P g C ⁻¹ | A |
| PC2 | Diatom C | g C m ⁻³ | S |
| GRPC2 | Grazing of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| SEPC2 | Settling of diatom C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.9 PP3, Cyanobacteria P, g P m⁻³

$$\frac{dPP3}{dt} = UPPP3 + UPLDOPP3 - GRPP3 - DEPP3 - SEPP3 - BUOYP3 \quad (3.63)$$

Where:

| Process | Comment | Unit |
|----------|--|-------------------------------------|
| UPPP3 | Uptake of PO ₄ into cyanobacteria P | g P m ⁻³ d ⁻¹ |
| UPLDOPP3 | Cyanobacteria uptake of LDOP | g P m ⁻³ d ⁻¹ |
| GRPP3 | Grazing of cyanobacteria P | g P m ⁻³ d ⁻¹ |
| DEPP3 | Death of cyanobacteria P | g P m ⁻³ d ⁻¹ |
| SEPP3 | Settling of cyanobacteria P | g P m ⁻³ d ⁻¹ |
| BUOYP3 | Upward movement cyanobacteria | g P m ⁻³ d ⁻¹ |

UPPP3: Uptake of PO₄ into cyanobacteria P, g P m⁻³ d⁻¹

$$UPPP3 = \text{MIN}(upo3, ppma * PRPC3) \quad (3.64)$$

UPLDOPP3: Cyanobacteria uptake of LDOP, g P m⁻³ d⁻¹

$$UPLDOPP3 = \text{IF } PO4 < 0.001 \text{ THEN} \\ \text{IF } \left(ppma - \frac{PP3}{PC3} \right) < \text{epsi}, 0, \text{maxupip} * pda3 * 0.1 * \frac{LDOP}{(LDOP + hupp * pdb3 * 0.1)} * PC3) \\ \text{ELSE } 0 \quad (3.65)$$

GRPP3: Grazing of cyanobacteria P, g P m⁻³ d⁻¹

$$GRPP3 = pp3pc3 * GRPC3 \quad (3.66)$$

DEPP3: Death of cyanobacteria P, g P m⁻³ d⁻¹

$$DEPP3 = pp3pc3 * DEPC3 \quad (3.67)$$

SEPP3: Settling of cyanobacteria P, g P m⁻³ d⁻¹

$$SEPP3 = pp3pc3 * SEPC3 \quad (3.68)$$

BUOYP3: Upward movement of PP3, g P m⁻³ d⁻¹

$$BUOYP3 = pp3pc3 * BUOYC3 \quad (3.69)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|-------------------------------------|--------|
| ppma | Max. intracellular algae P | g P g C ⁻¹ | C |
| upo3 | Potential PO ₄ uptake by cyanobacteria | g P m ⁻³ d ⁻¹ | A |
| epsi | Small value | n.u. | C |
| maxupip | Max. PO ₄ uptake by flagellates during P | g P g C ⁻¹ | C |
| LDOP | Labile DOP | g P m ⁻³ | S |
| pda3 | Ratio, nutrient uptake cyanobacteria:flagellates | n.u. | A |
| pdb3 | Halfsaturation conc. Cyanobacteria:flagellates | n.u. | A |
| pp3pc3 | Cyanobacteria P:C ratio | g P g C ⁻¹ | A |
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| PP3 | Cyanobacteria P | g C m ⁻³ | S |
| GRPC3 | Grazing of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| DEPC3 | Death of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| SEPC3 | Settling of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| BUOYC3 | Upward movement of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.10 PSi2, Diatom Si, g Si m⁻³

$$\frac{dPSi2}{dt} = UPSi2 - GRPSi2 - DEPSi2 - SEPSi2 \quad (3.70)$$

Where:

| Process | Comment | Unit |
|---------|------------------------------|--------------------------------------|
| UPSi2 | Uptake of Si into diatoms Si | g Si m ⁻³ d ⁻¹ |
| GRPSi2 | Grazing of diatoms Si | g Si m ⁻³ d ⁻¹ |
| DEPSi2 | Death of diatoms Si | g Si m ⁻³ d ⁻¹ |
| SEPSi2 | Settling of diatoms Si | g Si m ⁻³ d ⁻¹ |

UPSi2: Uptake of Si into diatom Si, g Si m⁻³ d⁻¹

$$UPSi2 = MIN(upsi, psma * PRPC2) \quad (3.71)$$

GRPSi2: Grazing of diatomSiP, g Si m⁻³ d⁻¹

$$GRPSi2 = psi2pc2 * GRPC2 \quad (3.72)$$

DEPSi2: Death of diatom Si, g Si m⁻³ d⁻¹

$$DEPSi2 = psi2pc2 * DEPC2 \quad (3.73)$$

SEPSi2: Settling of diatom Si, g Si m⁻³ d⁻¹

$$SEPSi2 = psi2pc2 * SEPC2 \quad (3.74)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|-------------------------------|--------------------------------------|--------|
| psma | Max. intracellular algae Si | g Si g C ⁻¹ | C |
| usi2 | potential Si uptake by diatom | g Si m ⁻³ d ⁻¹ | A |
| psi2pc2 | Diatom Si:C ration | Si P g C ⁻¹ | A |
| PC2 | Diatom C | g C m ⁻³ | S |
| GRPC2 | Grazing of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| SEPC2 | Settling of diatom C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.11 CH, Chlorophyll, g m⁻³

$$\frac{dCH}{dt} = PRCH - DECH - SECH - BUOYCH - GRCH \quad (3.75)$$

Where:

| Process | Comment | Unit |
|---------|--|---------------------------------------|
| PRCH | Net production phytoplankton chlorophyll | g Chl m ⁻³ d ⁻¹ |
| SECH | Settling of phytoplankton chlorophyll | g Chl m ⁻³ d ⁻¹ |
| DECH | Death of phytoplankton chlorophyll | g Chl m ⁻³ d ⁻¹ |
| BUOYCH | Upward movement of CH | g Chl m ⁻³ d ⁻¹ |
| GRCH | Zooplankton grazing on CH | g Chl m ⁻³ d ⁻¹ |

PRCH: Net production phytoplankton chlorophyll, g Chl m⁻³ d⁻¹

The production of chlorophyll

$$PRCH = PRPC1 * \frac{chmi}{ik1} * e^{chma+myn1} + PRPC2 * \frac{chmi}{ik2} * e^{chma+myn2} + PRPC3 * \frac{chmi}{ik1} * e^{chma+myn3} \quad (3.76)$$

SECH: Settling of phytoplankton chlorophyll, g Chl m⁻³ d⁻¹

$$SECH = \frac{CH}{PC1 + PC2 + PC3} * (SEPC1 + SEPC2 + SEPC3) \quad (3.77)$$

DECH: Death of phytoplankton chlorophyll, g Chl m⁻³ d⁻¹

$$DECH = \frac{CH}{PC1 + PC2 + PC3} * (DEPC1 + DEPC2 + DEPC3) \quad (3.78)$$

GRCH: ZC Grazing on CH, g Chl m⁻³ d⁻¹

$$GRCH = \frac{CH}{PC1 + PC2 + PC3} * (GRPC1 + GRPC2 + GRPC3) \quad (3.79)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|--|--|--------|
| PC1 | Flagellate C | g C m ⁻³ | S |
| PC2 | Diatom C | g C m ⁻³ | S |
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| chmi | Min. chlorophyll-a production | mol photon ⁻¹ m ⁻² d ⁻¹ | C |
| chma | Max. chlorophyll-a producti | mol photon ⁻¹ m ⁻² d ⁻¹ | C |
| myn1 | Nitrogen function Flagellate | n.u. | A |
| myn2 | Nitrogen function Diatom | n.u. | A |
| myn2 | Nitrogen function Cyanobacteria | n.u. | A |
| ik1 | Light saturation temp. corrected, PC1, PC3 | mol photon m ⁻² d ⁻¹ | A |
| lk2 | Light saturation temp. corrected, PC2 | mol photon m ⁻² d ⁻¹ | A |
| PRPC1 | Net production of flagellate C | g C m ⁻³ d ⁻¹ | P |
| SEPC1 | Sedimentation of flagellate C | g C m ⁻³ d ⁻¹ | P |
| DEPC1 | Death of flagellate C | g C m ⁻³ d ⁻¹ | P |
| GRPC1 | Grazing of flagellate C | g C m ⁻³ d ⁻¹ | P |
| PRPC2 | Net production of diatom C | g C m ⁻³ d ⁻¹ | P |
| SEPC2 | Sedimentation of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| GRPC2 | Grazing of diatom C | g C m ⁻³ d ⁻¹ | P |
| PRPC3 | Net production of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| SEPC3 | Sedimentation of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| DEPC3 | Death of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| GRPC3 | Grazing of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.12 ZC, zooplankton, g C m⁻³

$$\frac{dZC}{dt} = PRZC - DEZC \quad (3.80)$$

Where:

| Process | Comment | Unit |
|---------|--------------------------------------|-------------------------------------|
| PRZC | Net production of zooplankton carbon | g C m ⁻³ d ⁻¹ |
| DEZC | Death of zooplankton carbon | g C m ⁻³ d ⁻¹ |

PRZC: Production of zooplankton carbon, g C m⁻³ d⁻¹

$$PRZC = vefo * (GRPC1 + GRPC2 + GRPC3) \quad (3.81)$$

DEZC: Death of zooplankton carbon, g C m⁻³ d⁻¹

$$DEZC = kdz * zc^2 + kdzb * zc \quad (3.82)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|-----------------------------------|--------------------------------------|--------|
| vefo | Zooplankton growth efficiency | g C g C ⁻¹ | C |
| kdz | Zooplankton death rate 2nd order, | m ³ (g C*d) ⁻¹ | C |
| kdzb | Zooplankton death rate 1st order | d ⁻¹ | C |
| GRPC1 | Grazing of flagellate C | g C m ⁻³ d ⁻¹ | P |
| GRPC2 | Grazing of diatom C | g C m ⁻³ d ⁻¹ | P |
| GRPC3 | Grazing of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.13 DC, Detritus C, g C m⁻³

$$\frac{dDC}{dt} = EKZC + DEZC + DEPC2DC - REDC - deDCw - SREDC - SEDC \quad (3.83)$$

Where:

| Process | Comment | Unit |
|---------|---|-------------------------------------|
| EKZC | Excretion by zooplankton carbon | g C m ⁻³ d ⁻¹ |
| DEZC | Death of zooplankton carbon | g C m ⁻³ d ⁻¹ |
| DEPC2DC | Death phytoplankton to detritus carbon | g C m ⁻³ d ⁻¹ |
| REDC | DO mineralisation of detritus carbon | g C m ⁻³ d ⁻¹ |
| deDCw | DC anaerobic respiration with NO ₃ | g C m ⁻³ d ⁻¹ |
| SREDC | DC anaerobic oxidation with SO ₄ | g C m ⁻³ d ⁻¹ |
| SEDC | Settling of detritus carbon | g C m ⁻³ d ⁻¹ |

EKZC: Excretion by zooplankton carbon, g C m⁻³ d⁻¹

$$EKZC = (1 - vefo - refo) * (GRPC1 + GRPC2 + GRPC3) \quad (3.84)$$

DEZC: Death of zooplankton carbon, g C m⁻³ d⁻¹

DEZC: see processes for zooplankton C, ZC, Equation (3.82)

DEPC2DC: Death phytoplankton to detritus carbon, g C m⁻³ d⁻¹

$$DEPC2DC = (1 - vm - vp - vn) * (DEPC1 + DEPC2 + DEPC3) \quad (3.85)$$

REDC: DO mineralisation of detritus carbon, g C m⁻³ d⁻¹

$$EDC = kmdm * DC * tere^{T-20} * \frac{DO^{ndo2}}{MAX(0.01, DO^{ndo2} + mdo2)} \quad (3.86)$$

deDCw: DC respiration with NO₃, g C m⁻³ d⁻¹

$$deDCw = (DENW * vn3 + ANAMOX * 0.429) * \frac{DC}{DC + LDOC} \quad (3.87)$$

SREDC: DC oxidation with SO₄, g C m⁻³ d⁻¹

$$sredc = SRED * vso * \frac{dc}{dc + LDOC} \quad (3.88)$$

SEDC: Settling of detritus carbon, g C m⁻³ d⁻¹

$$SEDC = ksd * \frac{dc}{dz} \quad (3.89)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| vefo | Zooplankton growth effency | g C g C ⁻¹ | C |
| refo | Zooplankton, respiration | g C g C ⁻¹ | C |
| GRPC1 | Grazing of flagellate C | g C m ⁻³ d ⁻¹ | P |
| GRPC2 | Grazing of diatom C | g C m ⁻³ d ⁻¹ | P |
| GRPC3 | Grazing of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| DEPC1 | Death of flagellate C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC3 | Death of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| vm | Fraction of PC mineralised at PC death | n.u. | C |
| vp | Fraction of PC to CDOC-N&P at PC death | n.u. | C |
| vn | Fraction of PC to LDOC-N&P at PC death | n.u. | C |
| kmdm | DC mineralisation rate at 20 ° C | d ⁻¹ | C |
| tere | Θ in Arrhenius function, DC mineralisation | n.u. | C |
| T | Temperature | °C | F |
| DO | Oxygen | g O ₂ m ⁻³ | S |
| ndo2 | DC & LDOC:Coefficient, DO mineralisation | n.u. | C |
| mdo2 | DO half-saturation constant, DC & LDOC mineralisation | n.u | C |
| DENW | Denitrificaion in water using DC+LDOC | g N m ⁻³ d ⁻¹ | P |
| vn3 | C:N ratio denitrification | g C g N ⁻¹ | C |
| ANAMOX | Anammox, NO ₃ +NH ₄ → N ₂ | g N m ⁻³ d ⁻¹ | P |

| Name | Comment | Unit | Type*) |
|------|--|-------------------------------------|--------|
| LDOC | Labile DOC | g C m ⁻³ | S |
| SRED | SO ₄ Respiration of DC+LDOC | g S m ⁻³ d ⁻¹ | P |
| vso | C:S ratio C mineralisation SO ₄ to H ₂ S | g C g S ⁻¹ | C |
| ksd | Sedimentation rate detritus | m d ⁻¹ | A |
| dz | Height of actual water layer in model | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.14 DN, Detritus N, g N m⁻³

$$\frac{dDN}{dt} = EKZN + DEZN + DEPN2DN - REDN - SREDN - deDNw - SEDN \quad (3.90)$$

Where:

| Process | Comment | Unit |
|---------|---|-------------------------------------|
| EKZN | Excretion by zooplankton N | g N m ⁻³ d ⁻¹ |
| DEZN | Death of zooplankton N | g N m ⁻³ d ⁻¹ |
| DEPN2DN | Death phytoplankton to detritus N | g N m ⁻³ d ⁻¹ |
| REDN | DO mineralisation of detritus N to NH ₄ | g N m ⁻³ d ⁻¹ |
| deDNw | Anaerobic respiration of DN with NO ₃ to NH ₄ | g N m ⁻³ d ⁻¹ |
| SREDN | Anaerobic oxidation of DN with SO ₄ to NH ₄ | g N m ⁻³ d ⁻¹ |
| SEDN | Settling of detritus N | g N m ⁻³ d ⁻¹ |

EKZN: Excretion by zooplankton N, g N m⁻³ d⁻¹

$$EKZN = (1 - vefo - refo) * (GRPN1 + GRPN2 + GRPN3) \quad (3.91)$$

DEZN Death of zooplankton N, g N m⁻³ d⁻¹

$$DEZN = vzn * DEZC \quad (3.92)$$

DEPN2DN: Death phytoplankton to detritus N, g N m⁻³ d⁻¹

$$DEPN2DN = (1 - vm - vp - vn) * (DEPN1 + DEPN2 + DEPN3) \quad (3.93)$$

REDN: DO mineralisation of detritus N, g N m⁻³ d⁻¹

$$REDN = kmdn * dndc * REDC \quad (3.94)$$

deDNw: DN respiration with NO₃, g N m⁻³ d⁻¹

$$deDNw = deDCw * dndc \quad (3.95)$$

SREDN: DN oxidation with SO₄, g N m⁻³ d⁻¹

$$SREDN = SREDC * dndc \quad (3.96)$$

SEDN: Settling of detritus N, g N m⁻³ d⁻¹

$$SEDN = SEDC * dndc \quad (3.97)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|--|-------------------------------------|--------|
| vefo | Zooplankton growth effency | g C g C ⁻¹ | C |
| refo | Zooplankton, respiration | g C g C ⁻¹ | C |
| GRPN1 | Grazing of flagellate N | g N m ⁻³ d ⁻¹ | P |
| GRPN2 | Grazing of diatom N | g N m ⁻³ d ⁻¹ | P |
| GRPN3 | Grazing of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| DEPN1 | Death of flagellate N | g N m ⁻³ d ⁻¹ | P |
| DEPN2 | Death of diatom N | g N m ⁻³ d ⁻¹ | P |
| DEPN3 | Death of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| vzn | N:C ratio Zooplankton | g N g C ⁻¹ | C |
| vm | Fraction of PC mineralised at PC death | n.u. | C |
| vp | Fraction of PC to CDOC-N&P at PC death | n.u. | C |
| vn | Fraction of PC to LDOC-N&P at PC death | n.u. | C |
| kmdn | Factor N mineralisation of DN | n.u. | C |
| dndc | N:C ration, detritus | g N g C ⁻¹ | A |
| REDC | DO mineralisation of detritus carbon | g C m ⁻³ d ⁻¹ | P |

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------------------|--------|
| deDCw | Anaerobic DC respiration with NO ₃ | g C m ⁻³ d ⁻¹ | P |
| SREDC | Anaerobic DC oxidation with SO ₄ | g C m ⁻³ d ⁻¹ | P |
| SEDC | Settling of detritus carbon | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.15 DP, Detritus P, g P m⁻³

$$\frac{dDP}{dt} = EKZP + DEZP + DEPP2DP - REDP - SREDP - deDPw - SEDP \quad (3.98)$$

Where:

| Process | Comment | Unit |
|----------|---|-------------------------------------|
| EKZP | Excretion by zooplankton P | g P m ⁻³ d ⁻¹ |
| DEZP | Death of zooplankton P | g P m ⁻³ d ⁻¹ |
| DEPP2DNP | Death phytoplankton to detritus P | g P m ⁻³ d ⁻¹ |
| REDP | DO mineralisation of detritus P to PO ₄ | g P m ⁻³ d ⁻¹ |
| deDPw | Anaerobic respiration of DP with NO ₃ to PO ₄ | g P m ⁻³ d ⁻¹ |
| SREDP | Anaerobic oxidation of DP with SO ₄ to PO ₄ | g P m ⁻³ d ⁻¹ |
| SEDP | Settling of detritus P | g P m ⁻³ d ⁻¹ |

EKZP: Excretion by zooplankton P, g P m⁻³ d⁻¹

$$EKZP = (1 - vefo - refo) * (GRPP1 + GRPP2 + GRPP3) \quad (3.99)$$

DEZP: Death of zooplankton P, g P m⁻³ d⁻¹

$$DEZP = v zp * DEZC \quad (3.100)$$

DEPP2DP: Death phytoplankton to detritus P, g P m⁻³ d⁻¹

$$DEPP2DP = (1 - vm - vp - vn) * (DEPP1 + DEPP2 + DEPP3) \quad (3.101)$$

REDP: DO mineralisation of detritus P, g P m⁻³ d⁻¹

$$REDP = kmdp * dpdc * REDC \tag{3.102}$$

deDPw: DP respiration with NO₃, g P m⁻³ d⁻¹

$$deDPw = deDCw * dpdc \tag{3.103}$$

SREDP: DP oxidation with SO₄, g P m⁻³ d⁻¹

$$SREDP = SREDC * dpdc \tag{3.104}$$

SEDP: Settling of detritus P, g P m⁻³ d⁻¹

$$SEDP = SEDC * dpdc \tag{3.105}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------------------|--------|
| vefo | Zooplankton growth effency | g C g C ⁻¹ | C |
| refo | Zooplankton, respiration | g C g C ⁻¹ | C |
| GRPP1 | Grazing of flagellate P | g P m ⁻³ d ⁻¹ | P |
| GRPP2 | Grazing of diatom P | g P m ⁻³ d ⁻¹ | P |
| GRPP3 | Grazing of cyanobacteria P | g P m ⁻³ d ⁻¹ | P |
| DEPP1 | Death of flagellate P | g P m ⁻³ d ⁻¹ | P |
| DEPP2 | Death of diatom P | g P m ⁻³ d ⁻¹ | P |
| DEPP3 | Death of cyanobacteria P | g P m ⁻³ d ⁻¹ | P |
| vzp | P:C ratio Zooplankton | g P g C ⁻¹ | C |
| vm | Fraction of PC mineralised at PC death | n.u. | C |
| vp | Fraction of PC to CDOC-N&P at PC death | n.u. | C |
| vn | Fraction of PC to LDOC-N&P at PC death | n.u. | C |
| kmdp | Factor P mineralisation of DP | n.u. | C |
| dpdc | P:C ration, detritus | g P g C ⁻¹ | A |
| REDC | DO mineralisation of detritus carbon | g C m ⁻³ d ⁻¹ | P |
| deDCw | Anaerobic DC respiration with NO ₃ | g C m ⁻³ d ⁻¹ | P |

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------------------|--------|
| SREDC | Anaerobic DC oxidation with SO ₄ | g C m ⁻³ d ⁻¹ | P |
| SEDC | Settling of detritus carbon | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.16 DSi, Detritus Si, g Si m⁻³

$$\frac{dDSi}{dt} = DEPSi2 + GRPSi2 - REDSi - SEDSi \quad (3.106)$$

Where:

| Process | Comment | Unit |
|---------|--|--------------------------------------|
| DEPSi2 | Death phytoplankton Si to detritus Si | g Si m ⁻³ d ⁻¹ |
| GRPSi2 | Grazing of Diatom Si | g Si m ⁻³ d ⁻¹ |
| REDSi | DO mineralisation of detritus Si to Si | g Si m ⁻³ d ⁻¹ |
| SEDSi | Settling of detritus Si | g Si m ⁻³ d ⁻¹ |

DEPSi2: Death phytoplankton Si to detritus Si, g Si m⁻³ d⁻¹

$$DEPSi2 = DEPC2 * psi2pc2 \quad (3.107)$$

GRPSi2: Grazing of Diatom Si, g Si m⁻³ d⁻¹

$$GRSi2 = GRPC2 * psi2pc2 \quad (3.108)$$

REDSi: DO mineralisation of detritus Si to Si, g Si m⁻³ d⁻¹

$$REDSi = REDC * dsidc * km ds \quad (3.109)$$

SEDSi: Settling of detritus Si, g Si m⁻³ d⁻¹

$$SEDSi = SEDC * dsidc \quad (3.110)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|--------------------------------------|-------------------------------------|--------|
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| psi2pc2 | Si:C ration, Diatom | g Si g C ⁻¹ | A |
| GRPC2 | Grazing of diatom C | g C m ⁻³ d ⁻¹ | P |
| REDC | DO mineralisation of detritus carbon | g C m ⁻³ d ⁻¹ | P |
| dsidc | Si:C ration, detritus | g Si g C ⁻¹ | A |
| kmds | Factor Si mineralisation of DSi | n.u. | C |
| SEDC | Settling of detritus carbon | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.17 NH₄, Total ammonia, g N m⁻³

$$\begin{aligned} \frac{dNH_4}{dt} = & REDN + REZN + deDNw + SREDN + reLDON + deLDON \\ & + sreLDON + feunh4m3 + fsnb + NH_4dep + DEPN2NH \\ & - RNIT - ANAMOX - UPNH1 - UPNH2 - UPNH3 \end{aligned} \quad (3.111)$$

Where:

| Process | Comment | Unit |
|---------------------|--|-------------------------------------|
| REDN | DN→NH ₄ via DO oxidation of DC | g N m ⁻³ d ⁻¹ |
| REZN | Respiration of zooplankton nitrogen | g N m ⁻³ d ⁻¹ |
| deDNw | DN→NH ₄ denitrification mineralisation of DC | g N m ⁻³ d ⁻¹ |
| SREDN | DN →NH ₄ via SO ₄ mineralisation of DC | g N m ⁻³ d ⁻¹ |
| reLDON | LDON→NH ₄ via DO oxidation of LDOC | g N m ⁻³ d ⁻¹ |
| deLDON | LDON →NH ₄ via denitrification mineralisation of LDOC | g N m ⁻³ d ⁻¹ |
| sreLDON | LDON →NH ₄ via SO ₄ mineralisation of LDOC | g N m ⁻³ d ⁻¹ |
| feunh4m3 | NH ₄ flux between sediment pore water and water | g N m ⁻³ d ⁻¹ |
| fsnb | Mineralisation of newly settled organic N | g N m ⁻³ d ⁻¹ |
| NH ₄ dep | Atmospheric NH ₄ deposition | g N m ⁻³ d ⁻¹ |
| DEPN2NH | Fraction of DEPN1-3 to NH ₄ | g N m ⁻³ d ⁻¹ |

| Process | Comment | Unit |
|---------|---|---|
| RNIT | Nitrification in water column | $\text{g N m}^{-3} \text{d}^{-1}$ |
| ANAMOX | Anammox, $\text{NO}_3 + \text{NH}_4 \rightarrow \text{N}_2$ | $\text{g NH}_4\text{-N m}^{-3} \text{d}^{-1}$ |
| UPNH1 | NH_4 uptake by flagellate | $\text{g N m}^{-3} \text{d}^{-1}$ |
| UPNH2 | NH_4 uptake by diatom | $\text{g N m}^{-3} \text{d}^{-1}$ |
| UPNH3 | NH_4 uptake by cyanobacteria | $\text{g N m}^{-3} \text{d}^{-1}$ |

REDN: NH_4 production via mineralisation of DC, DN & DP with DO, $\text{g N m}^{-3} \text{d}^{-1}$

Please see under DN, Equation (3.94)

REZN: Respiration of zooplankton nitrogen, $\text{g N m}^{-3} \text{d}^{-1}$

$$REZN = (\text{GRPN1} + \text{GRPN2} + \text{GRPN3}) * \text{vzn} - \text{EKZN} \quad (3.112)$$

Where: **EKZN: Excretion of N by zooplankton**

$$EKZN = (1 - \text{vefo} - \text{refo}) * (\text{GRPN1} + \text{GRPN2} + \text{GRPN3}) \quad (3.113)$$

deDNw: NH_4 production via denitrification (NO_3 mineralisation) of DC, DN & DP, $\text{g N m}^{-3} \text{d}^{-1}$

Please see under DN, Equation (3.95)

SREDN: NH_4 production via anaerobic SO_4 mineralisation of DC, DN & DP, $\text{g N m}^{-3} \text{d}^{-1}$

Please see under DN, Equation (3.96)

reLDON: NH_4 production via mineralisation of LDOC, LDON & LDOP with DO, $\text{g N m}^{-3} \text{d}^{-1}$

Please see under LDON (Section 3.3.27).

deLDON: NH_4 production via denitrification, (mineralisation) of LDOC, LDON & LDOP, $\text{g N m}^{-3} \text{d}^{-1}$

Please see under LDON (Section 3.3.27).

sreLDON: NH_4 production via anaerobic SO_4 mineralisation of LDOC, LDON & LDOP, $\text{g N m}^{-3} \text{d}^{-1}$

Please see under LDON (Section 3.3.27).

feunh4m3: NH4 flux between sediment pore water and water, g N m⁻³ d⁻¹

$$feunh4m3 = feunh4/dz \tag{3.114}$$

fsnb: Mineralisation of newly settled organic N, g N m⁻³ d⁻¹

$$fsnb = krsn0 * (SEPN1 - BUOYN1 + SEPN2 + SEPN3 - BUOYN3 + SEDN - depoC * knim/dz) * tetn^{T-20} \tag{3.115}$$

NH4depo: Atmospheric N deposition as NH4 to surface layer, g N m⁻³ d⁻¹

$$NH4depo = NHdepo/dz \tag{3.116}$$

RNIT: Nitrification in water column, g N m⁻³ d⁻¹

$$RNIT = knitw * tnit^{T-20} * sqdo * \frac{NH4}{NH4 + hmt} * NH4 \tag{3.117}$$

ANAMOX: Anammox, NO3+NH4 → N2, , g NH4-N m⁻³ d⁻¹

ANAMOX = IF DO < 0.32 THEN

$$kanam * tetn^{T-20} * \frac{NH4}{NH4 + hun4} * \frac{NO3}{NO3 + hun3} * \frac{DC + LDOC}{DC + LDOC + hudc1} \tag{3.118}$$

ELSE 0

UPNH1-3: NH4 uptake by phytoplankton, g N m⁻³ d⁻¹

UPNH1, UPNH2 & UPNH3 see under PN1 (Section 3.3.4), PN2 (Section 3.3.5) and PN3 (Section 3.3.6)

Where:

| Name | Comment | Unit | Type*) |
|-------|-------------------------------|-------------------------------------|--------|
| vzn | N to C ratio in zooplankton, | g N g C ⁻¹ | C |
| GRPN1 | Grazing of flagellate N | g N m ⁻³ d ⁻¹ | P |
| GRPN2 | Grazing of diatom N | g N m ⁻³ d ⁻¹ | P |
| GRPN3 | Grazing of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| EKZN | Excretion of N by zooplankton | g N m ⁻³ d ⁻¹ | P |
| vefo | Zooplankton growth effency | g C g C ⁻¹ | C |

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| refo | Zooplankton, respiration | g C g C ⁻¹ | C |
| feunh4 | Flux of NH ₄ from sediment to water | g N m ⁻² d ⁻¹ | P |
| SEPN1 | Settling of flagellate N | g N m ⁻³ d ⁻¹ | P |
| SEPN2 | Settling of diatom N | g N m ⁻³ d ⁻¹ | P |
| SEPN3 | Settling of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| BUOYN1 | Upward movement of flagellate N | g N m ⁻³ d ⁻¹ | P |
| BUOYN3 | Upward movement of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| depoC | Deposition of organic C to sediment | g C m ⁻² d ⁻¹ | P |
| knim | Sediment: N:C ratio of immobile N | g N g C ⁻¹ | C |
| tetn | Θ value in Arrhenius equation for N | n.u. | C |
| T | Temperature Deg. Celsius | °C | C |
| NHdep | Atmospheric N deposition | g N m ⁻² d ⁻¹ | F |
| knitw | Specific nitrification water at 20 C | d ⁻¹ | C |
| tnit | Θ value in Arrhenius equation for nitrification | n.u. | C |
| hmt | Halfsaturation NH ₄ nitrification | g N m ⁻³ d ⁻¹ | C |
| sqdo | DO function | n.u. | A |
| kanam | max anammox NO ₃ -N or NH ₄ -N consumption | g N m ⁻³ d ⁻¹ | C |
| NO3 | NO ₃ -N | g N m ⁻³ | S |
| hun4 | NH ₄ half saturation conc., anammox & thiodenitrification | g N m ⁻³ | C |
| hun3 | NO ₃ half saturation concentration, anammox | g N m ⁻³ | C |
| DC | Detrituc C | g C m ⁻³ | S |
| LDOC | Labile DOC | g C m ⁻³ | S |
| hudc1 | DC+LDC Half saturation concentration, anammox | g C m ⁻³ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.18 NO₃, Nitrate, g N m⁻³

$$\frac{dNO_3}{dt} = RNIT + feuno3m3 + NO3dep - DENW - DENW_s - ANAMOX - UPN31 - UPN32 - UPN33 \quad (3.119)$$

Where:

| Process | Comment | Unit |
|-------------------|---|--|
| RNIT | Nitrification in water column | g N m ⁻³ d ⁻¹ |
| feuno3m3 | Flux of NO ₃ between water & sediment | g N m ⁻³ d ⁻¹ |
| NO3dep | Atmospheric deposition of NO ₃ at the water surface | g N m ⁻³ d ⁻¹ |
| DENW | Denirification water | g N m ⁻³ d ⁻¹ |
| DENW _s | Thiodenitification, 4NO ₃ +3H ₂ S → 2N ₂ +3SO ₄ | g N m ⁻³ d ⁻¹ |
| ANAMOX | Anammox, NO ₃ +NH ₄ → N ₂ | g NO ₃ -N m ⁻³ d ⁻¹ |
| UPN31 | NO ₃ uptake by flagellate | g N m ⁻³ d ⁻¹ |
| UPN32 | NO ₃ uptake by diatom | g N m ⁻³ d ⁻¹ |
| UPN32 | NO ₃ uptake by cyanobacteria | g N m ⁻³ d ⁻¹ |

RNIT: Nitrification in water column, g N m⁻³ d⁻¹

Please see under NH₄, Equation (3.117)

feuno3m3: Flux of NO₃ between water & sediment, g N m⁻³ d⁻¹

$$feuno3m3 = feuno3/dz \quad (3.120)$$

NO3dep: Atmospheric deposition of NO₃ at the water surface, g N m⁻³ d⁻¹

$$NO3dep = NO3depo/dz \quad (3.121)$$

DENW: Denirification water, g N m⁻³ d⁻¹

denw = IF DO < mdo3 THEN

$$kdenw * tetn^{T-20} * \frac{ksb + 0.1}{DC + LDOC} * \frac{NO3}{NO3 + hun3} * \frac{1}{DC + LDOC + hudc} \quad (3.122)$$

ELSE 0

DENWS: Thiodenitification, $4\text{NO}_3+3\text{H}_2\text{S}\rightarrow 2\text{N}_2+3\text{SO}_4$, $\text{g N m}^{-3} \text{d}^{-1}$

$DENW_s = \text{IF } DO < 0.32 \text{ THEN}$

$$kdenw * tetn^{T-20} * \frac{H2S}{H2S + hs1} * \frac{NO3}{NO3 + hun3} * \frac{hudc}{DC + LDOC + hudc} \quad (3.123)$$

ELSE 0

ANAMOX: Anammox, $\text{NO}_3+\text{NH}_4 \rightarrow \text{N}_2$, $\text{g NO}_3\text{-N m}^{-3} \text{d}^{-1}$

$ANAMOX = \text{IF } DO < 0.32 \text{ THEN}$

$$kanam * tetn^{T-20} * \frac{NH4}{NH4 + hun4} * \frac{NO3}{NO3 + hun3} * \frac{DC + LDOC}{DC + LDOC + hudc1} \quad (3.124)$$

ELSE 0

UPN31, UPN32, UPN33: NO_3 uptake flagellates, diatoms and cyanobacteria, $\text{g N m}^{-3} \text{d}^{-1}$

Please see under PN1 (Section 3.3.4), PN2 (Section 3.3.5) and PN3 (Section 3.3.6)

Where:

| Name | Comment | Unit | Type*) |
|---------|--|-----------------------------------|--------|
| feuno3 | Nitrate flux between sediment and water column | $\text{g N m}^{-2} \text{d}^{-1}$ | P |
| NO3depo | Atmospheric $\text{NO}_3\text{-N}$ deposition to water surface | $\text{g N m}^{-2} \text{d}^{-1}$ | F |
| mdo3 | DO limit for denitrification in water column & half saturation concentration in sqdo | $\text{g O}_2 \text{m}^{-3}$ | C |
| Kdenw | Max. denitrification at 20 °C in water column | $\text{g N m}^{-3} \text{d}^{-1}$ | C |
| ksb | Half saturation DO conc. for denitrification | $\text{g O}_2 \text{m}^{-3}$ | C |
| tetn | Θ in Arrhenius function, denitrifications temperature dependency | n.u | C |
| hun3 | NO_3 half saturation concentration for denitrification | g N m^{-3} | C |
| hudc | DC+LDOC Half saturation concentration for SO_4 reduction & denitrification | g C m^{-3} | C |
| DC | Detritus C | g C m^{-3} | S |
| LDOC | Labile fraction of DOC | g C m^{-3} | S |
| tetn | Θ value in Arrhenius equation for N | n.u. | C |

| Name | Comment | Unit | Type*) |
|-------|--|---------------------|--------|
| T | Temperature | °C | F |
| H2S | H ₂ S-S | g S m ⁻³ | S |
| hs1 | H ₂ S half saturation thiodenitrification | g S m ⁻³ | C |
| NH4 | NH ₄ -N | g N m ⁻³ | S |
| hun4 | NH ₄ half saturation conc., anammox & thiodenitrification | g N m ⁻³ | C |
| hudc1 | DC+LDC Half saturation concentration, anammox | g C m ⁻³ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.19 H₂S, Hydrogen Sulphide, g S m⁻³

$$\frac{dH2S}{dt} = SRED + fwsh2sm3 - SOXI - SOXI_N \tag{3.125}$$

Where:

| Process | Comment | Unit |
|-------------------|--|-------------------------------------|
| SRED | Anaerobic SO ₄ Respiration of DC+LDOC | g S m ⁻³ d ⁻¹ |
| fwsh2sm3 | H ₂ S flux from sediment to water | g S m ⁻³ d ⁻¹ |
| SOXI | Oxidation of H ₂ S | g S m ⁻³ d ⁻¹ |
| SOXI _N | SO ₄ production by thiodenitrification, 4NO ₃ +3H ₂ S→2N ₂ +3SO ₄ | g S m ⁻³ d ⁻¹ |

SRED: SO₄ Respiration of DC+LDOC, g S m⁻³ d⁻¹

SRED = IF DO < 0.5 THEN

$$ksc * fsa * ts4r^{(T-20)} * \frac{hun3}{NO3 + hun3} * \frac{DC + LDOC}{DC + LDOC + hudc} \tag{3.126}$$

ELSE 0

fwsh2sm3: H₂S flux from sediment to water, g S m⁻³ d⁻¹

$$fwsh2sm3 = fwh2s/dz \tag{3.127}$$

SOXI: Oxidation of H₂S, g S m⁻³ d⁻¹

$$SOXI = kse * ksf^{(T-20)} * H2S * sqdo \quad (3.128)$$

SOXI_N: SO₄ production by thiodenitrification, 4NO₃+3H₂S-->2N₂+3SO₄, g S m⁻³ d⁻¹

$$SOXI_N = DENW_s * 1.429 \quad (3.129)$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|--|-------------------------------------|--------|
| Ksc | Max. anoxic DC +LDOC respiration rate with SO ₄ | g S m ⁻³ d ⁻¹ | C |
| Fsa | Salinity function for reduction of SO ₄ to H ₂ S | n.u. | A |
| hun3 | NO ₃ half saturation concentration for denitrification | g N m ⁻³ | C |
| Hudc | DC+LDOC Half saturation concentration for SO ₄ reduction & denitrification | g C m ⁻³ | C |
| DC | Detritus C | g C m ⁻³ | S |
| LDOC | Labile fraction of DOC | g C m ⁻³ | S |
| ts4r | Θ value in Arrhenius function for SO ₄ reductions temperature dependency. | n.u. | C |
| fwh2s | Flux of reduced H ₂ S equivalents from sediment to water | g S m ⁻² d ⁻¹ | P |
| Kse | Max. specific oxidation rate of H ₂ S, 20 deg. °C | d ⁻¹ | C |
| Ksf | Θ value in Arrhenius function for SO ₄ oxidations temperature dependency. | n.u. | C |
| Sqdo | DO function | n.u. | A |
| DENW _s | Thiodenitrification. 4NO ₃ +3H ₂ S→2N ₂ +3SO ₄ | g N m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.20 IP, Phosphate (PO₄-P), g P m⁻³

$$\begin{aligned} \frac{dIP}{dt} = & REDP + REZP + reLDOP + deLDOP + sreLDOP + deDPw + SREDP \\ & + DEPP2IP + fspb + fsipm3 + Pdep - UPPP1 - UPPP2 \\ & - UPPP3 \end{aligned} \quad (3.130)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| REDP | DP→PO ₄ via DO oxidation of DC | g P m ⁻³ d ⁻¹ |
| REZP | Respiration of zooplankton P | g P m ⁻³ d ⁻¹ |
| reLDOP | LDOP→PO ₄ via DO oxidation of LDOC | g P m ⁻³ d ⁻¹ |
| deLDOP | LDOP →PO ₄ via denitrification mineralisation of LDOC | g P m ⁻³ d ⁻¹ |
| sreLDOP | LDOP →PO ₄ via SO ₄ mineralisation of LDOC | g P m ⁻³ d ⁻¹ |
| deDPw | DP→PO ₄ denitrification mineralisation of DC | g P m ⁻³ d ⁻¹ |
| SREDP | DP →PO ₄ via SO ₄ mineralisation of DC | g P m ⁻³ d ⁻¹ |
| DEPP2IP | Dead Plankton P to PO ₄ | g P m ⁻³ d ⁻¹ |
| fspb | Mineralisation of newly settled organic P | g P m ⁻³ d ⁻¹ |
| fsipm3 | PO ₄ flux between sediment pore water and water | g P m ⁻³ d ⁻¹ |
| Pdep | Atmospheric P deposition | g P m ⁻³ d ⁻¹ |
| UPPP1 | PO ₄ uptake by flagellates | g P m ⁻³ d ⁻¹ |
| UPPP2 | PO ₄ uptake by Diatoms | g P m ⁻³ d ⁻¹ |
| UPPP3 | PO ₄ uptake by cyanobacteria | g P m ⁻³ d ⁻¹ |

REDP: PO₄ production via mineralisation of DC, DN & DP with DO, g P m⁻³ d⁻¹

Please see under state variable DP, Section 3.3.15

REZP: Respiration of zooplankton phosphorus, g P m⁻³ d⁻¹

$$REZP = \text{MAX}(0, GRPP1 + GRPP2 + GRPP3 - PRZC * v_{zp} - EKZP) \quad (3.131)$$

reLDOP: PO₄ production via mineralisation of LDOC, LDON & LDOP with DO, g P m⁻³ d⁻¹

$$reLDOP = \frac{LDOP}{LDOC} * reLDOC \quad (3.132)$$

deLDOP: PO₄ production via denitrification (mineralisation) of LDOC, LDON & LDOP, g P m⁻³ d⁻¹

$$deLDOP = \frac{LDOP}{LDOC} * deLDOC \quad (3.133)$$

sreLDOP: PO₄ production via anaerobic SO₄ mineralisation of LDOC, LDON & LDOP, g P m⁻³ d⁻¹

$$sreLDOP = \frac{LDOP}{LDOC} * sreLDOC \quad (3.134)$$

deDPw: PO₄ production via anaerobic denitrification (mineralisation) of DC, DN & DP, g P m⁻³ d⁻¹

Please see under DP, Section 3.3.15

SREDP: PO₄ production via anaerobic SO₄ mineralisation of DC, DN & DP, g P m⁻³ d⁻¹

Please see under DP, Section 3.3.15

DEPP2IP: Dead Plankton P to PO₄ , g P m⁻³ d⁻¹

$$DEPP2IP = (DEPP1 + DEPP2 + DEPP3) * vm \quad (3.135)$$

fspb: Mineralisation of newly settled organic P, g P m⁻³ d⁻¹

$$fspb = krsp0 * (SEPP1 - BUOYP1 + SEPP2 + SEPP3 - BUOYP3 + SEDP) * tetp^{T-20} \quad (3.136)$$

fsipm3: PO₄ flux between sediment pore water and water, g P m⁻³ d⁻¹

$$fsipm3 = fsip/dz \quad (3.137)$$

Pdep: Atmospheric P deposition as PO₄ to surface layer, g P m⁻³ d⁻¹

$$Pdep = Pdepo/dz \quad (3.138)$$

UPPP1-3: PO₄ uptake by flagellates, diatoms and cyanobacteria, g P m⁻³ d⁻¹

Please see under PP1 (Section 3.3.7), PP2 (Section 3.3.8), and PP3 (Section 3.3.9)

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------|--------|
| GRPP1 | Grazing of phytoplankton (Flagellate) P | $g P m^{-3} d^{-1}$ | P |
| GRPP2 | Grazing of phytoplankton (diatom) P | $g P m^{-3} d^{-1}$ | P |
| GRPP3 | Grazing of cyanobacteria P | $g P m^{-3} d^{-1}$ | P |
| PRZC | Net production of zooplankton C | $g C m^{-3} d^{-1}$ | P |
| vzp | P:C ratio in zooplankton | $g P g C^{-1}$ | C |
| EKZP | Excretion by zooplankton P | $g P m^{-3} d^{-1}$ | P |
| reLDOC | DO respiration LDOC | $g C m^{-3} d^{-1}$ | P |
| LDOP | Labile DOP | $g P m^{-3}$ | S |
| LDOC | Labile DOC | $g C m^{-3}$ | S |
| deLDOC | Anaerobic mineralisation of LDOC via denitrification | $g C m^{-3} d^{-1}$ | P |
| sreLDOC | Anaerobic mineralisation of LDOC via SO_4 reduction | $g C m^{-3} d^{-1}$ | P |
| DEPP1 | Death of flagellate P | $g P m^{-3} d^{-1}$ | P |
| DEPP2 | Death of diatom P | $g P m^{-3} d^{-1}$ | P |
| DEPP3 | Death of cyanobacteria P | $g P m^{-3} d^{-1}$ | P |
| vm | Fraction of PC mineralised at PC death | n.u. | C |
| krsp0 | Fraction of newly settled P to mineralisation | n.u. | C |
| SEPP1 | Sedimentation of flagellates P | $g P m^{-3} d^{-1}$ | P |
| SEPP2 | Sedimentation of diatom P | $g P m^{-3} d^{-1}$ | P |
| SEPP3 | Sedimentation of cyanobacteria P | $g P m^{-3} d^{-1}$ | P |
| BUOYP1 | Upward movement of flagellates P | $g P m^{-3} d^{-1}$ | P |
| BUOYP3 | Upward movement of cyanobacteria P | $g P m^{-3} d^{-1}$ | P |
| SEDP | Sedimentation detritus P | $g P m^{-3} d^{-1}$ | P |
| tetp | Θ value in Arrhenius equation for P | n.u. | C |
| T | Temperature Deg. Celsius | $^{\circ}C$ | C |
| fsip | PO_4 flux between pore water and water | $g P m^{-2} d^{-1}$ | P |
| Pdep | Atmospheric P deposition | $g P m^{-2} d^{-1}$ | F |
| dz | Height of actual water layer in model | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.21 IP, Phosphate (PO₄-P), g P m⁻³

$$\frac{dSi}{dt} = REDSi + RSSi2Si - UPSi2 \quad (3.139)$$

Where:

| Process | Comment | Unit |
|---------|---|--------------------------------------|
| REDSi | DO mineralisation of detritus DSi to Si | g Si m ⁻³ d ⁻¹ |
| RSSi2Si | Si release fom sediment | g Si m ⁻³ d ⁻¹ |
| UPSi2 | Uptake of Si into diatom Si | g Si m ⁻³ d ⁻¹ |

DEPSi2: Death phytoplankton Si to detritus Si, g Si m⁻³ d⁻¹

Please see under DSi, Section 3.3.16

RSSi2Si: Si release fom sediment, g Si m⁻³ d⁻¹

$$RSSi2Si = RSSi/dz \quad (3.140)$$

UPSi2: Uptake of Si into diatom Si, g Si Si m⁻³ d⁻¹

Please see under Psi2, Section 3.3.10

Where:

| Name | Comment | Unit | Type*) |
|------|---------------------------------------|--------------------------------------|--------|
| RSSi | Si release form sediment | g Si m ⁻² d ⁻¹ | P |
| dz | Height of actual water layer in model | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.22 DO, Oxygen, g O₂ m⁻³

$$\frac{dDO}{dt} = ODPC + REAR + REAR1 - ODDC - ODZC - SOXI2DO - RNIT2DO - reLDOC2DO - DEPC2DO - ODSC \quad (3.141)$$

Where:

| Process | Comment | Unit |
|-----------|--|--|
| ODPC | Net O ₂ production by phytoplankton | g O ₂ m ⁻³ d ⁻¹ |
| REAR | Reaeration | g O ₂ m ⁻³ d ⁻¹ |
| REAR1 | Reaeration shallow water | g O ₂ m ⁻³ d ⁻¹ |
| ODDC | DO consumption by mineralisation of DC | g O ₂ m ⁻³ d ⁻¹ |
| ODZC | Zooplankton respiration | g O ₂ m ⁻³ d ⁻¹ |
| SOXI2DO | DO consumption due to H ₂ S oxidation | g O ₂ m ⁻³ d ⁻¹ |
| RNIT2DO | DO consumption due to nitrification | g O ₂ m ⁻³ d ⁻¹ |
| reLDOC2DO | DO consumption by mineralisation of LDOC | g O ₂ m ⁻³ d ⁻¹ |
| DEPC2DO | DO consumption by mineralisation during PC death | g O ₂ m ⁻³ d ⁻¹ |
| ODSC | Sediment DO consumption | g O ₂ m ⁻³ d ⁻¹ |

ODPC: Net O₂ production/consumption by phytoplankton, g O₂ m⁻³d⁻¹

$$ODPC = (PRPC1 + PRPC2 + PRPC3) * v_o \quad (3.142)$$

REAR: Reaeration, surface layer only, g O₂ m⁻³d⁻¹

$$REAR = IF \text{ depth} < 5 \text{ THEN } 0 \text{ ELSE} \\ \left(3.93 * \frac{\sqrt{hcur}}{\text{depth}^{1.5}} + \frac{2.07 + 0.215 * wsp^{1.7}}{100 * 24} \right) * (csair - DO) / dz \quad (3.143)$$

REAR1: Reaeration, shallow water all layers, g O₂ m⁻³d⁻¹

$$REAR1 = IF \text{ depth} < 5 \text{ THEN} \\ \left(3.93 * \frac{\sqrt{hcur}}{\text{MAX}(0.1, \text{depth})^{1.5}} + \frac{2.07 + 0.215 * wsp^{1.7}}{100 * 24} \right) * (csair - DO) * dz / \text{depth} \\ \text{ELSE } 0 \quad (3.144)$$

ODDC: DO consumption by mineralisation of DC, g O₂ m⁻³d⁻¹

$$ODDC = REDC * v_o \quad (3.145)$$

ODZC: Zooplankton respiration, g O₂ m⁻³d⁻¹

$$ODZC = REZC * v_o \quad (3.146)$$

Where:

$$REZC = (GRPC1 + GRPC2 + GRPC3) * refo \quad (3.147)$$

SOXI2DO: DO consumption due to H₂S oxidation, g O₂ m⁻³d⁻¹

$$SOXI2DO = SOXI * vsh \quad (3.148)$$

RNIT2DO: DO consumption due to nitrification, g O₂ m⁻³d⁻¹

$$RNIT2DO = RNIT * vmh \quad (3.149)$$

reLDOC2DO: DO consumption by mineralisation of LDOC, g O₂ m⁻³d⁻¹

$$reLDOC2do = reLDOC * v_o \quad (3.150)$$

DEPC2DO: DO consumption by mineralisation during PC death, g O₂ m⁻³d⁻¹

$$dDEPC2DO = (DEPC1 + DEPC2 + DEPC3) * vm * v_o \quad (3.151)$$

ODSC: Sediment DO consumption, g O₂ m⁻³d⁻¹

From the sediment-water interface oxygen (DO) can penetrate into the sediment pore water by diffusion or actively being transported into the sediment by ventilation pumping and sediment mixing by the benthic fauna. Further microbenthic algae through photosynthesis can produce DO in the sediment-water interface. DO is consumed in the sediment by bacterial respiration and chemical oxidation of reduced substances (Fe⁺⁺, H₂S) resulting in the O₂ concentration becoming 0 (normally 0-2 cm) below the sediment surface. In the model this depth is defined as KDO2. Assuming the DO produced by the microbenthic algae is delivered to the water, the below differential equation can be set up assuming a steady state condition:

$$0 = -difo2 * \frac{d^2 O_2}{dy^2} + DOconsum \quad (3.152)$$

Where $0 < y < (KDO2)$, which by integration becomes:

$$\frac{d O_2}{dy} = \frac{DOconsum}{difo2} * y + a \quad (3.153)$$

Where a is a constant, which by using the border condition ($dO_2/dy=0$ at $y=KDO2$) can be defined as:

$$a = -\frac{DOconsum}{difo2} * KDO2 \Rightarrow \quad (3.154)$$

$$\frac{dO_2}{dy} = \frac{DOconsum}{difo2} * y - \frac{DOconsum}{difo2} * KDO2$$

Which by yet an integration gives:

$$O_2 = \frac{DOconsum}{2 * difo2} * y^2 - \frac{DOconsum}{difo2} * KDO2 * y + b \quad (3.155)$$

Where b is a constant, which by using the border condition ($O_2=0$ at $y=KDO2$) can be defined as:

$$b = \frac{DOconsum}{2 * difo2} * KDO2^2 \Rightarrow \quad (3.156)$$

$$O_2 = \frac{DOconsum}{2 * difo2} * y^2 - \frac{DOconsum}{difo2} * KDO2 * y + \frac{DOconsum}{difo2} * KDO2^2$$

At the sediment surface $y=0$ the $O_2 = DO \Rightarrow$

$$KDO2 = \sqrt{2 * difo2 * \frac{DO}{DOconsum}} \Rightarrow \quad (3.157)$$

$$O_2 = \frac{DOconsum}{2 * difo2} * y^2 - \frac{DOconsum}{difo2} * \sqrt{2 * difo2 * \frac{DO}{DOconsum}} * y + 2 * DO$$

The flux of DO from the water into the sediment can be described using Fick's 1. Law at depth $y=0$

$$DOflux = -difo2 * \frac{dO_2}{dy} \quad (3.158)$$

$\frac{dO_2}{dy}$ is found by differentiation of the above expression for O_2 in the sediment and determine the flux of DO into the sediment at $Y=0$.

$$DOflux = -difo2 * \left(-\frac{DOconsum}{difo2} * \sqrt{2 * difo2 * \frac{DO}{DOconsum}} \right) \Rightarrow \quad (3.159)$$

$$DOflux = \sqrt{2 * DO * difo2 * DOconsum} \Rightarrow$$

The DO consumption in the model is the sum of bacterial respiration ($reKDO2$), nitrification ($rsnit$) and a flux of reduced substances from the under laying sediment ($fsh2s$) to the layer with O_2 . All the mentioned DO consuming processes has the unit ($g\ m^{-2}d^{-1}$) and therefore has to be divided with the DO penetration ($KDO2$). A conversion factor for $O_2:N$ of 4.57 $g\ O_2 : g\ NH_4-N$ is used and a conversion factor for $O_2:S$ of 2 $g\ O_2 : H_2S-S$ is used.

The diffusion or rather transport of oxygen into the sediment is dependent of the activity of the benthic infauna. Their activity is linked to the DO concentration, at low DO (below 2 g m⁻³) the activity will decrease caused by increased mortality. The constant *difO2* is therefore multiplied by an oxygen function (1+sqdo).

The final equation for (ODSC, g O₂ m⁻³d⁻¹) in the template therefore becomes:

$$ODSC = \begin{cases} \text{IF } KDO2 > 0.001 \text{ THEN} \\ \sqrt{2 * difO2 * (1 + sqdo) * DO * \frac{fsh2s * 2 + rsnit * 4.57 + reKDO2}{KDO2}} * \frac{1}{dz} \\ \text{ELSE } 0 \end{cases} \quad (3.160)$$

Where:

| Name | Comment | Unit | Type* |
|-------|--|-------------------------------------|-------|
| PRPC1 | Production flagellate carbon | g C m ⁻³ d ⁻¹ | P |
| PRPC2 | Production diatom carbon | g C m ⁻³ d ⁻¹ | P |
| PRPC3 | Production cyanobacteria carbon | g C m ⁻³ d ⁻¹ | P |
| vo | O ₂ :C ratio for Production & respiration | g O ₂ g C ⁻¹ | C |
| hcur | Horizontal current | m s ⁻¹ | F |
| wsp | Wind speed, 10 m above sea | m s ⁻¹ | F |
| CSAIR | O ₂ saturation in water, relative to PSU & temp. | g O ₂ m ⁻³ | A |
| DO | Oxygen in water | g O ₂ m ⁻³ | S |
| depth | Depth of water column | m | F |
| dz | Height of actual water layer | m | F |
| REDC | Respiration detritus | g C m ⁻³ d ⁻¹ | P |
| REZC | Respiration zooplankton | g C m ⁻³ d ⁻¹ | P |
| GRPC1 | Grazing of phytoplankton (Flagellate) carbon | g C m ⁻³ d ⁻¹ | P |
| GRPC2 | Grazing of phytoplankton (diatom) carbon | g C m ⁻³ d ⁻¹ | P |
| GRPC3 | Grazing of cyanobacteria carbon | g C m ⁻³ d ⁻¹ | P |
| refo | Zooplankton, respiration | g C g C ⁻¹ | C |
| SOXI | H ₂ S oxidation to SO ₄ | g S m ⁻² d ⁻¹ | P |
| vsh | O ₂ :S ratio for oxidation of H ₂ S to SO ₄ | g O ₂ g S ⁻¹ | C |
| RNIT | Nitrification | g N m ⁻³ d ⁻¹ | P |

| Name | Comment | Unit | Type*) |
|----------|---|--|--------|
| vnh | O ₂ :N ratio nitrification | g O ₂ g N ⁻¹ | C |
| reLDOC | Respiration LDOC | g C m ⁻³ d ⁻¹ | P |
| DEPC1 | Death of flagellate C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC3 | Death of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| vm | Fraction of DEPC1-3 respired at once | n.u. | C |
| difO2 | Diffusion of O ₂ in sediment | m ² s ⁻¹ | C |
| DOconsum | Sediment O ₂ consumption, layer (0-KDO2) | g O ₂ m ⁻³ d ⁻¹ | |
| y | Depth below sediment surface | m | |
| sqdo | DO dependend auxiliary | n.u. | A |
| fsh2s | H ₂ S flux from under laying anoxic sediment layer | g S m ⁻² d ⁻¹ | P |
| rsnit | Nitrification in sediment in oxic sediment layer | g N m ⁻² d ⁻¹ | P |
| reKDO2 | Respiration in oxic sediment layer | g O m ⁻² d ⁻¹ | P1 |
| KDO2 | Oxic layer in sediment | m | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.23 CDOC, Coloured refractory DOC, g C m⁻³

$$\frac{dCDOC}{dt} = depc2CDOC - phoxCDOC \tag{3.161}$$

Where:

| Process | Comment | Unit |
|-----------|------------------------------------|-------------------------------------|
| depc2CDOC | Fraction of depc to CDOC | g C m ⁻³ d ⁻¹ |
| phoxCDOC | UV Photo oxidation of CDOC to LDOC | g C m ⁻³ d ⁻¹ |

depc2CDOC: Fraction of depc to CDOC, g C m⁻³ d⁻¹

$$depc2CDOC = (DEPC1 + DEPC2 + DEPC3) * vp \tag{3.162}$$

phoxCDOC: UV photo oxidation of CDOC to LDOC, g C m⁻³ d⁻¹

$$phoxCDOC = CDOC * doc_{maxde} * rd * doc_{monod} \quad (3.163)$$

Where:

| Name | Comment | Unit | Type*) |
|----------------------|--|-------------------------------------|--------|
| DEPC1 | Death of flagellate C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC3 | Death of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| vp | Fraction of DEPC, DEPN & DEPP to CDOC, CDON & CDOP | n.u. | C |
| doc _{maxde} | Max relative photo oxidation rate | d ⁻¹ | C |
| rd | Relative daylength, f(latitude, day,month,year) | n.u. | A |
| doc _{monod} | UV radiation Monod relation for photo oxidation | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.24 CDON, Coloured refractory DON, g N m⁻³

$$\frac{dCDON}{dt} = depc2CDON - phoxCDON \quad (3.164)$$

Where:

| Process | Comment | Unit |
|-----------|------------------------------------|-------------------------------------|
| depc2CDON | Fraction of depn to CDON | g N m ⁻³ d ⁻¹ |
| phoxCDON | UV Photo oxidation of CDON to LDON | g N m ⁻³ d ⁻¹ |

Depc2CDON: Fraction of depn to CDON, g N m⁻³ d⁻¹

$$depc2CDON = (DEPN1 + DEPN2 + DEPN3) * vp \quad (3.165)$$

phoxCDON: UV photo oxidation of CDON to LDON, g N m⁻³ d⁻¹

$$phoxCDON = CDON * doc_{maxde} * rd * doc_{monod} \quad (3.166)$$

Where:

| Name | Comment | Unit | Type*) |
|----------------------|--|-------------------------------------|--------|
| DEPN1 | Death of flagellate N | g N m ⁻³ d ⁻¹ | P |
| DEPN2 | Death of diatom N | g N m ⁻³ d ⁻¹ | P |
| DEPN3 | Death of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| vp | Fraction of depc, depn, depp to CDOC, CDON, CDOP | n.u. | C |
| rd | Relative daylength, f(latitude, day,month,year) | n.u. | A |
| doc _{maxde} | Max relative photo oxidation rate | d ⁻¹ | C |
| doc _{monod} | UV radiation Monod relation for photo oxidation | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.25 CDOP, Coloured refractory DOP, g P m⁻³

$$\frac{dCDOP}{dt} = depc2CDOP - phoxCDOP \tag{3.167}$$

Where:

| Process | Comment | Unit |
|-----------|------------------------------------|-------------------------------------|
| depc2CDOP | Fraction of depp to CDOP | g P m ⁻³ d ⁻¹ |
| phoxCDOP | UV Photo oxidation of CDOP to LDOP | g P m ⁻³ d ⁻¹ |

depc2CDOP: Fraction of depp to CDOP, g P m⁻³ d⁻¹

$$depc2CDOP = (DEPP1 + DEPP2 + DEPP3) * vp \tag{3.168}$$

phoxCDOP: UV photo oxidation of CDOP to LDOP, g P m⁻³ d⁻¹

$$phoxCDOP = CDOP * doc_{maxde} * rd * doc_{monod} \tag{3.169}$$

Where:

| Name | Comment | Unit | Type*) |
|----------------------|--|-------------------------------------|--------|
| DEPP1 | Death of flagellate P | g P m ⁻³ d ⁻¹ | P |
| DEPP2 | Death of diatom P | g P m ⁻³ d ⁻¹ | P |
| DEPP3 | Death of cyanobacteria P | g P m ⁻³ d ⁻¹ | P |
| vp | Fraction of depc, depn, depp to CDOC, CDON, CDOP | n.u. | C |
| rd | Relative daylength, f(latitude, day,month,year) | n.u. | A |
| doc _{maxde} | Max relative photo oxidation rate | d ⁻¹ | C |
| doc _{monod} | UV radiation Monod relation for photo oxidation | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.26 LDOC, Labile DOC, g C m⁻³

$$\frac{dLDOC}{dt} = phoxCDOC + depc2LDOC - reLDOC - deLDOC - sreLDOC \quad (3.170)$$

Where:

| Process | Comment | Unit |
|-----------|---|-------------------------------------|
| phoxCDOC | UV Photo oxidation of CDOC to LDOC | g C m ⁻³ d ⁻¹ |
| depc2LDOC | Fraction of depc to LDOC | g C m ⁻³ d ⁻¹ |
| reLDOC | Aerobic respiration of LDOC using O ₂ | g C m ⁻³ d ⁻¹ |
| deLDOC | Anaerobic respiration of LDOC using NO ₃ | g C m ⁻³ d ⁻¹ |
| sreLDOC | Anaerobic respiration of LDOC using SO ₄ | g C m ⁻³ d ⁻¹ |

phoxCDOC: UV photo oxidation of CDOC to LDOC, g C m⁻³ d⁻¹

See under CDOC, Section 3.3.23.

depc2CDPC: Fraction of depc to LDOC, g C m⁻³ d⁻¹

$$depc2CDOC = (DEPC1 + DEPC2 + DEPC3) * vn \quad (3.171)$$

reLDOC: Aerobic respiration of LDOC using O₂, g C m⁻³ d⁻¹

$$reLDOC = LDOC * kmLC * tere^{T-20} * \frac{DO^{ndo2}}{DO^{ndo2} + mdo2} \quad (3.172)$$

deLDOC: Anaerobic respiration of LDOC using NO₃, g C m⁻³ d⁻¹

$$deLDOC = \frac{LDOC}{DC + LDOC} * (DENW * vn3 + ANAMOX * 0.429) \quad (3.173)$$

sreLDOC: Anaerobic respiration of LDOC using SO₄, g C m⁻³ d⁻¹

$$sreLDOC = \frac{LDOC}{DC + LDOC} * SRED * vso \quad (3.174)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------------------|--------|
| DEPC1 | Death of flagellate C | g C m ⁻³ d ⁻¹ | P |
| DEPC2 | Death of diatom C | g C m ⁻³ d ⁻¹ | P |
| DEPC3 | Death of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| vn | Fraction of depc, depn, depp to LDOC, LDON, LDOP | n.u. | C |
| tere | Θ in Arrhenius function, DC mineralisation | n.u. | C |
| ndo2 | DC & LDOC:Coefficient, DO mineralisation | n.u. | C |
| mdo2 | DO half-saturation constant, DC & LDOC mineralisation | n.u. | C |
| kmLC | Specific mineralisation rate of LDOC at 20 °C | d ⁻¹ | C |
| DC | Detritus C | g C m ⁻³ | S |
| DENW | Denitrification in water | g N m ⁻³ d ⁻¹ | P |
| vn3 | C:N ratio denitrification | g C g N ⁻¹ | C |
| sred | Anoxic C mineralisation via SO ₄ →H ₂ S | g S m ⁻³ d ⁻¹ | P |
| vso | C:S ratio, SO ₄ respiration | g C g S ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.27 LDON, Labile DON, g N m^{-3}

$$\frac{dLDON}{dt} = phoxCDON + depc2LDON - reLDON - deLDON - sreLDON \quad (3.175)$$

Where:

| Process | Comment | Unit |
|-----------|---|-----------------------------------|
| phoxCDON | UV Photo oxidation of CDON to LDON | $\text{g N m}^{-3} \text{d}^{-1}$ |
| depc2LDON | Fraction of depn to LDON | $\text{g N m}^{-3} \text{d}^{-1}$ |
| reLDON | Aerobic respiration of LDON using O_2 | $\text{g N m}^{-3} \text{d}^{-1}$ |
| deLDON | Anaerobic respiration of LDON using NO_3 | $\text{g N m}^{-3} \text{d}^{-1}$ |
| sreLDON | Anaerobic respiration of LDON using SO_4 | $\text{g N m}^{-3} \text{d}^{-1}$ |

phoxCDON: UV photo oxidation of CDON to LDON, $\text{g N m}^{-3} \text{d}^{-1}$

See under CDON, Section 3.3.24.

depc2LDON: Fraction of depn to LDON, $\text{g N m}^{-3} \text{d}^{-1}$

$$depc2CDON = (DEPN1 + DEPN2 + DEPN3) * vn \quad (3.176)$$

reLDON: Aerobic respiration of LDON using O_2 , $\text{g N m}^{-3} \text{d}^{-1}$

$$reLDON = \frac{LDON}{LDOC} * reLDOC \quad (3.177)$$

deLDON: Anaerobic respiration of LDON using NO_3 , $\text{g N m}^{-3} \text{d}^{-1}$

$$deLDON = \frac{LDON}{LDOC} * deLDOC \quad (3.178)$$

sreLDON: Anaerobic respiration of LDON using SO_4 , $\text{g N m}^{-3} \text{d}^{-1}$

$$sreLDON = \frac{LDON}{LDOC} * sreLDOC \quad (3.179)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|-------------------------------------|--------|
| DEPN1 | Death of flagellate P | g N m ⁻³ d ⁻¹ | P |
| DEPN2 | Death of diatom N | g N m ⁻³ d ⁻¹ | P |
| DEPN3 | Death of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| vn | Fraction of depc, depn, depp to LDOC, LDON, LDOP | n.u. | C |
| reLDOC | Aerobic respiration of LDOC using O ₂ | g C m ⁻³ d ⁻¹ | P |
| deLDOC | Anaerobic respiration of LDOC using NO ₃ | g C m ⁻³ d ⁻¹ | P |
| sreLDOC | Anaerobic respiration of LDOC using SO ₄ | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.3.28 LDOP, Labile DOP, g P m⁻³

$$\frac{dLDOP}{dt} = phoxCDOP + depp2LDOP - UPLDOPP3 - reLDOP - deLDOP - sreLDOP \tag{3.180}$$

Where:

| Process | Comment | Unit |
|-----------|---|-------------------------------------|
| phoxCDOP | UV Photo oxidation of CDOP to LDOP | g P m ⁻³ d ⁻¹ |
| depp2LDOP | Fraction of depp to LDOP | g P m ⁻³ d ⁻¹ |
| UPLDOPP3 | Cyanobacteria uptake of LDOP | g P m ⁻³ d ⁻¹ |
| reLDOP | Aerobic respiration of LDOP using O ₂ | g P m ⁻³ d ⁻¹ |
| deLDOP | Anaerobic respiration of LDOP using NO ₃ | g P m ⁻³ d ⁻¹ |
| sreLDOP | Anaerobic respiration of LDOP using SO ₄ | g P m ⁻³ d ⁻¹ |

phoxCDOP: UV photo oxidation of CDOP to LDOP, g P m⁻³ d⁻¹

See under CDOP, Section 3.3.25.

depp2LDOP: Fraction of depp to LDOP, g P m⁻³ d⁻¹

$$depp2CDOP = depp * vn \tag{3.181}$$

UPLDOPP3: Cyanobacteria uptake of LDOP, g P m⁻³ d⁻¹

Please see under state variable PP3, Section 3.3.9.

reLDOP: Aerobic respiration of LDOP using O₂, g P m⁻³ d⁻¹

$$reLDOP = \frac{LDOP}{LDOC} * reLDOC \quad (3.182)$$

deLDOP: Anaerobic respiration of LDOP using NO₃, g P m⁻³ d⁻¹

$$deLDOP = \frac{LDOP}{LDOC} * deLDOC \quad (3.183)$$

sreLDOP: Anaerobic respiration of LDOP using SO₄, g P m⁻³ d⁻¹

$$sreLDOP = \frac{LDOP}{LDOC} * sreLDOC \quad (3.184)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|-------------------------------------|--------|
| DEPP1 | Death of flagellate P | g N m ⁻³ d ⁻¹ | P |
| DEPP2 | Death of diatom P | g N m ⁻³ d ⁻¹ | P |
| DEPP3 | Death of cyanobacteria P | g N m ⁻³ d ⁻¹ | P |
| Vn | Fraction of depc, depn, depp to LDOC, LDON, LDOP | n.u. | C |
| reLDOC | Aerobic respiration of LDOC using O ₂ | g C m ⁻³ d ⁻¹ | P |
| deLDOC | Anaerobic respiration of LDOC using NO ₃ | g C m ⁻³ d ⁻¹ | P |
| sreLDOC | Anaerobic respiration of LDOC using SO ₄ | g C m ⁻³ d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4 Differential Equation Sediment State Variables

3.4.1 S*S*_i, Sediment, bio-available Silicate, g Si m²³

$$\frac{dSSi}{dt} = SESi - RSSi \tag{3.185}$$

Where:

| Process | Comment | Unit |
|---------|------------------------------------|-------------------------------------|
| SESi | Deposition of Diatom & Detritus Si | g Si m ² d ⁻¹ |
| RSSi | Flux of Si from sediment | g Si m ² d ⁻¹ |

SESi: Deposition of Diatom & Detritus Si, g Si m²d⁻¹

$$SESi = (SEPSi2 + SEDSi) * dz \tag{3.186}$$

RSSi: Flux of Si from sediment, g Si m²d⁻¹

$$RSSi = krss * trss^{T-20} * \frac{SSi}{SSi + hss1} * MAX(1, MIN(2, \frac{0.005}{KDOX})) \tag{3.187}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|--------------------------------------|--------|
| SEPSi2 | Sedimentation of diatom Si | g Si m ⁻³ d ⁻¹ | P |
| SEDSi | Sedimentation of detritus Si | g Si m ⁻³ d ⁻¹ | P |
| dz | Height of actual water layer | M | F |
| krss | Max Si release rate from sediment at 20 °C | g Si m ⁻² d ⁻¹ | C |
| trss | Θ value in Arrhenius temperature function, Si | n.u. | C |
| T | Temperature | °C | F |
| hss1 | Half saturation constant for S <i>S</i> _i | g Si m ² | C |
| KDOX | NO ₃ penetration depth in sediment | M | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.2 KDOX, depth of NO₃ penetration in sediment, m

KDOX is the NO₃ penetration into the sediment. NO₃ is denitrified in the anoxic part of the sediment and therefore normally only penetrate 0-10 cm into the sediment. Normally, DO only penetrates a few mm into the sediment and therefore KDO2 (the DO penetration) will be smaller than KDOX. In a simulation, a situation may occur where KDOX is smaller than KDO2, which at least in theory may happen in nature. In this case, the increase in KDOX is set to a fixed fraction of the difference between KDO2 and KDOX.

$$\frac{dKDOX}{dt} = dkdox \quad (3.188)$$

Where:

| Process | Comment | Unit |
|---------|--------------------------------------|-------------------|
| Dkdox | change oxidised layer sediment, KDOX | m d ⁻¹ |

Change oxidised layer sediment, KDOX:

```

dkdox =IF KDOX<KDO2
THEN
(KDOX – KDO2) * kkdox
ELSE
dkdox_no3
    
```

(3.189)

Where:

| Name | Comment | Unit | Type* |
|-----------|--|-------------------|-------|
| KDO2 | DO penetration into sediment | m | S |
| kkdox | NO ₃ penetration rate constant into sediment, KDOX<KDO2 | d ⁻¹ | C |
| dkdox_no3 | NO ₃ penetration rate sediment, analytical solution | m d ⁻¹ | P1 |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.3 KDO2, DO penetration in sediment, m

$$\frac{dKDO2}{dt} = dkdo2 \tag{3.190}$$

Where:

| Process | Comment | Unit |
|---------|--------------------------------------|-------------------|
| dkdo2 | change in DO penetration in sediment | m d ⁻¹ |

dkdo2: Change in DO penetration in sediment, m d⁻¹

```

dkdo2 =IF (kds-KDO2)<epsi
THEN
MIN(kdo2i - KDO2) * kkdo2,0)
ELSE
(kdo2i-KDO2)*kkdo2
    
```

(3.191)

Where:

| Name | Comment | Unit | Type*) |
|-------|--|------|--------|
| KDO2 | DO penetration into sediment | m | S |
| kds | Depth of modelled sediment layer | m | C |
| epsi | Constant small value also used for PC nutrient uptake | n.u. | C |
| kdo2i | New steady state condition for KDO2, function of DO and respiration, analytical solution | m | P1 |
| kkdo2 | Rate constant for DO penetration into sediment | C | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.4 SOC, Sediment organic C, g C m⁻²

$$\frac{dSOC}{dt} = depoC - minSOC - rscim \quad (3.192)$$

Where:

| Process | Comment | Unit |
|---------|------------------------------|-------------------------------------|
| depoC | Deposition of C | g C m ⁻² d ⁻¹ |
| minSOC | Mineralisation of SOC | g C m ⁻² d ⁻¹ |
| Rscim | Burial of sediment organic C | g C m ⁻² d ⁻¹ |

Deposition of C on sediment surface:

$$depoC = (SEPC1 - BUOYC1 + SEP2 + SEPC3 - BUOYC3 + SEDC) * dz \quad (3.193)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| SEPC1 | Sedimentation of flagellate C | g C m ⁻³ d ⁻¹ | P |
| SEPC2 | Sedimentation of diatom C | g C m ⁻³ d ⁻¹ | P |
| SEPC3 | Sedimentation of cyanobacteria C | g C m ⁻³ d ⁻¹ | P |
| BUOYC1 | Flagellate upward movement | g C m ⁻³ d ⁻¹ | P |
| BUOYC3 | Cyanobacteria upward movement | g C m ⁻³ d ⁻¹ | P |
| SEDC | Sedimentation of detritus C (DC) to sediment | g C m ⁻³ d ⁻¹ | P |
| Dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

minSOC: Mineralisation of SOC, g C m⁻²d⁻¹

$$\text{minSOC} = \text{krsc1} * \text{SOC} * \text{tetn}^{T-20} + \text{fscb} * \text{dz} \quad (3.194)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------------------|--------|
| krsc1 | Specific mineralisation rate of SOC 20 °C | d ⁻¹ | C |
| tetn | Θ in Arrhenius temperature equation, SON mineralisation | n.u. | C |
| temp | Temperature | °C | F |
| fscb | Mineralisation of newly settled organic C | g C m ⁻³ d ⁻¹ | P1 |
| Dz | Height of actual layer= layer above sediment | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

rscim: Burial of sediment organic C, g C m⁻²d⁻¹

$$\text{rscim} = (\text{depoC} - \text{fscb} * \text{dz}) * \frac{\text{rsnim}}{\text{rson}} \quad (3.195)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------------------|--------|
| depoC | Deposition of C | g C m ⁻² d ⁻¹ | P |
| Fscb | Mineralisation of newly settled organic C | g C m ⁻³ d ⁻¹ | P1 |
| Rsnim | Burial of organic sediment N (SON), see SON (Section 3.4.5) | g N m ⁻² d ⁻¹ | P |
| Rson | Burial of organic sediment N (SON), see SON (Section 3.4.5) | g N m ⁻² d ⁻¹ | P |
| Dz | Height of actual layer= layer above sediment | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.5 SON, Bio-available organic N in sediment, g N m⁻²

$$\frac{dSON}{dt} = r_{son} - r_{sonnh} - r_{snim} \quad (3.196)$$

Where:

| Process | Comment | Unit |
|---------|---|-------------------------------------|
| Rson | Settling of organic N to SON | g N m ⁻² d ⁻¹ |
| Rsonnh | Mineralisation of SON to pore water NH ₄ | g N m ⁻² d ⁻¹ |
| Rsnim | Burial of organic sediment N (SON) | g N m ⁻² d ⁻¹ |

rson: Settling of organic N to SON

$$r_{son} = (SEPN1 - \mathbf{BUOYN1} + SEPN2 + SEPN3 - \mathbf{BUOYN3} + SEDN - f_{snb}) * dz \quad (3.197)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|-------------------------------------|--------|
| SEPN1 | Sedimentation of flagellate N | g N m ⁻³ d ⁻¹ | P |
| SEPN2 | Sedimentation of diatom N | g N m ⁻³ d ⁻¹ | P |
| SEPN3 | Sedimentation of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| BUOYN1 | Flagellate upward movement N | g N m ⁻³ d ⁻¹ | P |
| BUOYN3 | Cyanobacteria upward movement N | g N m ⁻³ d ⁻¹ | P |
| SEDN | Sedimentation of detritus N, (DN), see DN | g N m ⁻³ d ⁻¹ | P |
| Fsnb | Mineralisation of newly settled organic N on sed. surface | g N m ⁻³ d ⁻¹ | P1 |
| Dz | Height of actual layer= layer above sediment | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

rsonnh: Mineralisation of SON to pore water NH₄

$$rsonh = krsn1 * SON * tetn^{T-20} \tag{3.198}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-----------------|--------|
| krsn1 | Specific mineralisation rate of SON 20 °C | d ⁻¹ | C |
| Tetn | Θ in Arrhenius temperature equation, SON mineralisation | n.u. | C |
| T | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

rsnim: Burial of organic sediment N (SON), g N m⁻²d⁻¹

The mineralisation of SOC and SON is assumed to a function of the sediment SON:SOC ratio. At low SON:SOC ratios close to *knim* the mineralisation is assumed to small or close to 0. The fraction of organic N settled to the sediment (*rson*) to buried or immobilised is set to be *knim* multiplied with settled organic C to sediment. If N:C ration of the settled N is below *knim* all settled N (*rson*) is buried.

$$\begin{aligned}
 rsnim &= IF\ knim * (depoC - fscb * dz) < rson \\
 &THEN \\
 &knim * (depoC - fscb * dz) \\
 &ELSE \\
 &Rson
 \end{aligned} \tag{3.199}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------------------|--------|
| knim | Sediment: N:C ratio of immobile N | g N g C ⁻¹ | C |
| depoC | Deposition of C, see SOC | g C m ⁻² d ⁻¹ | P |
| fscb | Mineralisation of newly settled organic C | g C m ⁻³ d ⁻¹ | P1 |
| rson | Settling of organic N to SON | g N m ⁻² d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.6 SOP, Bio-available organic P in sediment, g P m⁻²

$$\frac{dSOP}{dt} = rsop - ropsip - rspim \quad (3.200)$$

Where:

| Process | Comment | Unit |
|---------|---|-------------------------------------|
| rsop | Settling of organic P to SOP | g P m ⁻² d ⁻¹ |
| ropsip | Mineralisation of SOP to pore water PO ₄ | g P m ⁻² d ⁻¹ |
| rspim | Burial – immobilisation of organic sediment P (SOP) | g P m ⁻² d ⁻¹ |

rsop: Settling of organic P to SOP, g P m⁻²d⁻¹

$$rsop = (SEPP1 - \mathbf{BUOYP1} + SEPP2 + SEPP3 - \mathbf{BUOYP3} + SEDP - fspb) * dz \quad (3.201)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|-------------------------------------|--------|
| SEPP1 | Sedimentation of flagellate P | g P m ⁻³ d ⁻¹ | P |
| SEPP2 | Sedimentation of diatom P | g P m ⁻³ d ⁻¹ | P |
| SEPP3 | Sedimentation of cyanobacteria P | g P m ⁻³ d ⁻¹ | P |
| BUOYP1 | Flagellate upward movement P | g P m ⁻³ d ⁻¹ | P |
| BUOYP3 | Cyanobacteria upward movement P | g P m ⁻³ d ⁻¹ | P |
| Fspb | Mineralisation of newly settled organic P on sed. Surface | g P m ⁻³ d ⁻¹ | P1 |
| Dz | Height of actual layer= layer above sediment | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

ropsip: Mineralisation of SOP to pore water PO₄, g P m⁻²d⁻¹

$$ropsip = krsp1 * SOP * tetp^{T-20} \tag{3.202}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-----------------|--------|
| krsp1 | Specific mineralisation rate of SOP 20 °C | d ⁻¹ | C |
| tetp | Θ in Arrhenius temperature equation, SOP mineralisation | n.u. | C |
| T | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

rspim: Burial and immobilisation of organic sediment P (SOP), g P m⁻²d⁻¹

P is able to be incorporated into hydroxyapatite (chalk) or adsorbed to reduced Fe (in fresh waters) or simply it is incorporated into inert organic C. The processes are not well known and therefore a fixed fraction of the settled organic P is immobilised. In sediments (typical marine) where the pool of H₂S (reduced substances) exceeds a fixed value of 0.01 g S m⁻² immobilisation to Fe⁺⁺ is reduced to 1/10.

The user may change the process if new information is available, or information on sediment type prescribes another formulation.

$$rspim = IF H2S > 0.01 THEN \frac{kpim}{10} * rsop ELSE kpim * rsop \tag{3.203}$$

Where:

| Name | Comment | Unit | Type*) |
|------|-----------------------------------|-------------------------------------|--------|
| knim | Sediment: P:C ratio of immobile P | g P g C ⁻¹ | C |
| rsop | Settling of organic P to SOP | g N m ⁻² d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.7 FESP, PO₄ adsorbed to oxidised iron in sediment, g P m⁻²

$$\frac{dFESP}{dt} = r_{fesip} \quad (3.204)$$

Where:

| Process | Comment | Unit |
|--------------------|---|-------------------------------------|
| r _{fesip} | Flux between pore water PO ₄ and iron-adsorbed P | g P m ⁻² d ⁻¹ |

r_{fesip}: Flux between pore water PO₄ and iron-adsorbed P, g P m⁻²d⁻¹

The exchange of PO₄-P between oxidised iron (Fe⁺⁺⁺) is calculated as a rate constant (*krap*) multiplied with the difference between a new steady state sorption of PO₄ to Fe⁺⁺⁺ (*FESP_f(KDOX)_∞*) and the last calculated pool of sorbed PO₄-P (*FESP_t*).

An approximation of *FESP_f(KDOX)_∞* is estimated using a Monod kinetic for PO₄ in the pore water combined with information of sediment iron content, dry matter sediment, density and finally multiplied with the oxidised layer (KDOX), which is a state variable in the model.

$$r_{fesip} = krap * (FESP_f(KDOX)_\infty - FESP_t) \Rightarrow$$

$$r_{fesip} = krap * (kfe * kfepo * \frac{SIPm3}{SIPm3 + khfe} * vf * dm * KDOX * 10^6 - FESP) \quad (3.205)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-------------------------|--------|
| krap | Rate constant for iron absorption – desorption of PO ₄ | d ⁻¹ | C |
| kfe | Oxidable iron content in sediment | g Fe g dw ⁻¹ | C |
| kfepo | Maximum Fe-P sorption capacity | g P g Fe ⁻¹ | C |
| SIPm3 | Pore water PO ₄ concentration | g P m ⁻³ | A |
| khfe | Half saturation constant, adsorption-desorption | g P m ⁻³ | C |
| vf | Sediment density | g WW cm ⁻³ | C |
| dm | Dry weight sediment | g DM g WW ⁻¹ | C |
| KDOX | Depth of NO ₃ penetration –oxidised layer with Fe ⁺⁺⁺ | m | S |
| FESP | Iron(Fe ⁺⁺⁺) bound PO ₄ in sediment | g P m ⁻² | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.8 SNH, Sediment pore water NH₄, g N m⁻²

$$\frac{dSNH}{dt} = rsonnh - rsnit - feunh4 \tag{3.206}$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| rsonnh | Mineralisation of SON to pore water NH ₄ | g N m ⁻² d ⁻¹ |
| rsnit | Nitrification of NH ₄ in sediment | g N m ⁻² d ⁻¹ |
| feunh4 | NH ₄ Flux between sediment pore water and water | g N m ⁻² d ⁻¹ |

rsonnh: mineralisation of SON to pore water NH₄, g N m⁻²d⁻¹

Please see under the state variable SON, Section 3.4.5.

rsnit: Nitrification of NH₄ in sediment, g N m⁻²d⁻¹

The nitrification of NH₄ in sediment pore water (*SNHm3*) is described as the product between a specific nitrification rate (*knit*), *SNHm3*, a Monod relation of *SNHm3* a Monod relation for DO (*sqdo*), the DO penetration in the sediment (*KDO2*) and a temperature relation.

$$rsnit = \text{IF } DO > 0 \text{ THEN } knit * SNHm3 * \frac{SNHm3}{SNHm3 + ksnh0} * sqdo * KDO2 * tetn^{T-20} \tag{3.207}$$

ELSE
0

Where:

| Name | Comment | Unit | Type*) |
|-------|---|---------------------|--------|
| Knit | Specific nitrification rate at 20 C in sediment | d ⁻¹ | C |
| SNHm3 | NH ₄ concentration in pore water | g N m ⁻³ | A |
| ksnh0 | NH ₄ half saturation concentration for nitrification | g N m ⁻³ | C |
| Sqdo | DO Mond function | n.u. | A |
| KDO2 | DO penetration in sediment | m | S |
| Tetn | Θ in Arrhenius temperature equation, SON mineralisation | n.u. | C |
| T | Temperature | °C | F |

- *) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

feunh4: NH₄ Flux between sediment pore water and water, g N m⁻²d⁻¹

The flux between pore water NH₄ (*SNHm3*) and NH₄ in the water (*NH4*) is described as a product of a vertical diffusion constant (*difnh*) and concentration difference divided with the the NO₃ penetration depth in the sediment (*KDOX*).

$$feunh4 = difnh * \frac{SNHm3 - NH4}{MIN(kds, KDOX)} \quad (3.208)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|--|--------------------------------|--------|
| difnh | Vertical diffusion for ammonia | m ² d ⁻¹ | C |
| SNHm3 | NH ₄ concentration in pore water | g N m ⁻³ | A |
| NH4 | NH ₄ concentration in water | g N m ⁻³ | C |
| kds | Depth of modelled sediment layer | m | C |
| KDOX | NO ₃ penetration in sediment-oxidised layer | m | S |

- *) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.9 SNO₃, NO₃ in sediment pore water, layer (0 - kdo2), g N m⁻²

$$\frac{dSNO3}{dt} = rsnit - rdenit - feuno3 \quad (3.209)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| rsnit | Nitrification of pore water NH ₄ in sediment | g N m ⁻² d ⁻¹ |
| Rdenit | Denitrification of NO ₃ in sediment | g N m ⁻² d ⁻¹ |
| feuno3 | NO ₃ Flux between sediment pore water and water | g N m ⁻² d ⁻¹ |

rsnit: Nitrification of pore water NH₄ in sediment, g N m⁻²d⁻¹

Please see state variable SNH, Section 3.4.8.

rdenit: Denitrification of NO₃ in sediment, g N m⁻²d⁻¹

rdenit is expressed as flux of NO₃ into the anoxic zone of the sediment where only diffusion and denitrification are the driving processes. An analytical solution is used assuming steady state conditions.

From the surface layer of the sediment with DO (KDO_2), NO₃ in the pore water with concentration SNO_3m_3 will penetrate deeper into the anoxic sediment layer while being denitrified to N₂. At a sudden depth below the surface ($KDOX$) NO₃ concentration will be 0. Assuming a constant denitrification and under steady state condition the NO₃ concentration (NO_3x) in the pore water at depth x below the surface layer with DO can be described with:

$$0 = -difno_3 * \frac{d^2NO_3x}{dx^2} + dnm_3 \text{ Where } 0 < x < (KDOX - KDO_2) \quad (3.210)$$

Which by integration becomes:

$$\frac{dNO_3x}{dx} = \frac{dnm_3}{difno_3} * x + a \quad (3.211)$$

Where a is a constant, which by using the border condition ($dNO_3x/dx=0$ at $x=(KDOX - KDO_2)$) can be defined as:

$$a = -\frac{dnm_3}{difno_3} * (KDOX - KDO_2) \Rightarrow \quad (3.212)$$

$$\frac{dNO_3x}{dx} = \frac{dnm_3}{difno_3} * (x - (KDOX - KDO_2))$$

Which by yet an integration gives:

$$NO_3x = \frac{dnm_3}{2 * difno_3} * x^2 * -\frac{dnm_3}{difno_3} * (KDOX - KDO_2) * x + b \quad (3.213)$$

Where b is a constant, which by using the border condition ($SNO_3m_3=0$ at $x=(KDOX - KDO_2)$) can be defined as:

$$b = \frac{dnm_3}{2 * difno_3} * (KDOX - KDO_2)^2 \Rightarrow \quad (3.214)$$

$$NO_3x = \frac{dnm_3}{2 * difno_3} * x^2 - \frac{dnm_3}{difno_3} * (KDOX - KDO_2) * x + \frac{dnm_3}{2 * difno_3} * (KDOX - KDO_2)^2$$

At depth $x=0$ in the anoxic zone (which is at depth KDO_2 below sediment surface) $NO_3x=SNO_3m_3$. \Rightarrow

$$(KDOX - KDO_2) = \sqrt{2 * difno_3 * \frac{SNO_3m_3}{dnm_3}} \Rightarrow \quad (3.215)$$

$$NO_3x = \frac{dnm_3}{2 * difno_3} * x^2 - \frac{dnm_3}{difno_3} * \sqrt{2 * difno_3 * \frac{SNO_3m_3}{dnm_3}} * x + 2 * SNO_3m_3$$

Assuming the flux of NO₃ into the anoxic sediment solely being created by denitrification the NO₃-flux=denitrification pr. m² d⁻¹ = rdenit.

Using Fick's 1. law for a flux at depth $x=0$.

$$r_{denit} = -d_{ifno3} * \frac{dNO_3x}{dx} \quad (3.216)$$

$\frac{dNO_3x}{dx}$ is found by differentiation of the above expression for NO_3x and determine the flux for $x=0$. The final equation used in the template:

$$r_{denit} = \sqrt{2 * d_{ifno3} * d_{nm3} * SNO_3m3} \quad (3.217)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-------------------------------------|--------|
| difno3 | Vertical diffusion for NO ₃ in sediment | m ² d ⁻¹ | C |
| dnm3 | denitrification in sediment, corrected for temperature | g N m ⁻³ d ⁻¹ | A |
| SNO3m3 | NO ₃ in pore water surface sediment, layer (0-kdo2) | g N m ⁻³ | A |
| KDO2 | DO penetration into the sediment | m | S |
| KDOX | NO ₃ penetration into the sediment | m | S |
| NO3x | NO ₃ concentration in pore water at depth x | g N m ⁻³ | |
| X | Depth below zone with DO | m | |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

feuno3: NO₃ Flux between sediment pore water and water, g N m⁻²d⁻¹

The flux between pore water NO₃ (SNO_3m3) and NO₃ in the water (NO_3) is described as a product of a vertical diffusion constant (d_{ifno3}) and concentration difference divided with the the DO penetration depth in the sediment ($KDO2$).

$$f_{euno3} = d_{ifno3} * \frac{SNO_3m3 - NO_3}{KDO2} \quad (3.218)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|--------------------------------|--------|
| difno3 | Vertical diffusion for NO ₃ in sediment | m ² d ⁻¹ | C |
| NO3 | NO ₃ concentration in water | g N m ⁻³ | S |
| SNO3m3 | NO ₃ in pore water surface sediment, layer (0-kdo2) | g N m ⁻³ | A |
| KDO2 | DO penetration into the sediment | m | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.10 SIP, PO₄ in sediment pore water, g P m⁻²

$$\frac{dSIP}{dt} = -rfesip + ropsip - fsip \tag{3.219}$$

Where:

| Process | Comment | Unit |
|---------|---|-------------------------------------|
| rfesip | Flux between pore water PO ₄ and iron-adsorbed P | g P m ⁻² d ⁻¹ |
| ropsip | Mineralisation of SOP to pore water PO ₄ | g P m ⁻² d ⁻¹ |
| fsip | Flux between PO ₄ in pore water and water above sediment | g P m ⁻² d ⁻¹ |

rfesip: Flux between pore water PO₄ and iron-adsorbed P, g P m⁻²d⁻¹

Please see under the stat variable FESP, sediment ion adsorbed PO₄, Section 3.4.7.

ropsip: Mineralisation of SOP to pore water PO₄, g P m⁻²d⁻¹

Please see under the stat variable SOP, bio-available organic P in sediment. Section 3.4.6.

fsip: Flux between PO₄ in pore water and water above sediment, g P m⁻²d⁻¹

The flux between pore water PO₄ (*SIPm3*) and PO₄ in the water (*IP*) is described as a product of a vertical diffusion constant (*kfip*) and concentration difference divided with the NO₃ penetration depth in the sediment (*KDOX*).

$$fsip = kfip * \frac{SIPm3 - IP}{KDOX} \tag{3.220}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|--|--------------------------------|--------|
| kfip | Vertical diffusion for PO ₃ in sediment | m ² d ⁻¹ | C |
| IP | PO ₄ concentration in water | g N m ⁻³ | S |
| SIPm3 | PO ₄ in pore water of the sediment | g N m ⁻³ | A |
| KDOX | NO ₃ penetration into the sediment | m | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.11 SH₂S, Reduced substances in sediment, g S m⁻²

$$\frac{dH_2S}{dt} = RSH_2S - fsh_2s - fwsh_2s \quad (3.221)$$

Where:

| Process | Comment | Unit |
|--------------------|---|-------------------------------------|
| RSH ₂ S | Sediment H ₂ S production in anoxic layer: mineralisation of SOC – denitrification | g S m ⁻² d ⁻¹ |
| fsh ₂ s | flux of SH ₂ S from reduced sediment (below KDOX) to oxidised sediment. | g S m ⁻² d ⁻¹ |
| fwsh ₂ | flux of reduced H ₂ S equivalents from sediment to water | g S m ⁻² d ⁻¹ |

RSH₂S: Sediment H₂S production in anoxic layer: mineralisation of SOC minus denitrification

The production of H₂S (reduced substances in sediment expressed as H₂S-S) is calculated as the mineralisation of SOC (minSOC) minus the a fraction of the minSOC being oxidised by DO in layer KDO₂ minus a fraction of minSOC being oxidised by NO₃ by denitrification in the anoxic zone penetrated by NO₃ (KDOX-KDO₂). A C:N ratio of 1.07 is used to convert denitrified NO₃-N to C, and a C:S ratio of 1.33 is used to convert mineralised C to S.

$$RSH_2S = \left(minSOC - \frac{erKDO_2}{v_o} - rdenit * 1.07 \right) * 1.33 \quad (3.222)$$

Where:

| Name | Comment | Unit | Type*) |
|--------------------|---|--|--------|
| minSOC | mineralisation SOC | g C m ⁻² d ⁻¹ | P |
| reKDO ₂ | Sediment mineralisation of SOC by DO, in layer KDO ₂ | g O ₂ m ⁻² d ⁻¹ | P1 |
| V _o | O ₂ : C ratio used in production & consumption processes | g O ₂ g C ⁻¹ | C |
| Rdenit | Denitrification in anoxic sediment layer | g N m ⁻² d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fsh2s: flux of SH2S from reduced sediment (below KDOX) to oxidised sediment, g S m⁻²d⁻¹

The flux of reduced substances expressed as H₂S-S from sediment below the oxidised zone of the sediment (below KDOX) is expressed as a diffusion constant multiplied by a H₂S concentration difference between the H₂S concentration (calculated from SH2S) and the H₂S concentration in the water above the sediment divided by the distance from the reduced zone in the sediment to the depth of DO penetration (KDO2).

$$\begin{aligned}
 & fsh2s = IF\ KDO2 > 0.001 \\
 & THEN \\
 & \quad \frac{SH2S}{(1 - dm) * vf * (kds - MAX(KDO2, (KDOX - KDO2)))} - H2S \quad (3.223) \\
 & difh2s * \frac{SH2S}{KDOX - KDO2} \\
 & ELSE\ 0
 \end{aligned}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|--------------------------------|--------|
| difh2s | Vertical diffusion of SH2S in sediment | m ² d ⁻¹ | C |
| SH2S | Reduced substances in sediment as H ₂ S | g S m ⁻² | S |
| vf | Sediment density | g WW cm ⁻³ | C |
| dm | Dry weight sediment | g DM g WW ⁻¹ | C |
| kds | Depth of modelled sediment layer | m | C |
| KDO2 | DO penetration into the sediment | m | S |
| KDOX | NO ₃ penetration into the sediment | m | S |
| H2S | Hydrogen sulphide in water above sediment (H ₂ S) | g S m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fwsh2s: flux of reduced H2S equivalents from sediment to water, g S m⁻² d⁻¹

In case the DO penetration in the sediment (KDO2) is below 1 mm the flux of H₂S go directly from the reduced zone of the sediment into the water above the sediment.

$$\begin{aligned}
 &fwsh2s = IF\ KDO2 \leq 0.001 \\
 &THEN \\
 &\quad difh2s * \frac{SH2S}{\frac{(1-dm)*vf*(kds-KDO2) - H2S}{KDOX}} \\
 &ELSE\ 0
 \end{aligned} \tag{3.224}$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|--|--------------------------------|--------|
| difh2s | Vertical diffusion of SH ₂ S in sediment | m ² d ⁻¹ | C |
| SH ₂ S | Reduced substances in sediment as H ₂ S | g S m ⁻² | S |
| Vf | Sediment density | g WW cm ⁻³ | C |
| Dm | Dry weight sediment | g DM g WW ⁻¹ | C |
| Kds | Depth of modelled sediment layer | m | C |
| KDO2 | DO penetration into the sediment | m | S |
| KDOX | NO ₃ penetration into the sediment | m | S |
| H ₂ S | Hydrogen sulphide in water above sediment (H ₂ S) | g S m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.12 SPIM, Immobilised sediment P, g P m⁻²

$$\frac{dSPIM}{dt} = rspim \tag{3.225}$$

Where:

| Process | Comment | Unit |
|---------|------------------------------|-------------------------------------|
| Rspim | Immobilisation of sediment P | g P m ⁻² d ⁻¹ |

rspim: Immobilisation of sediment P, g P m⁻²d⁻¹

The immobilisation of P in the sediment is set to be a constant fraction of the organic P settling to the sediment surface. In water bodies with permanently or semi permanent anoxia having a H₂S concentration above 0.01 g S m⁻³ the immobilisation is set to be 10% of

$$rspim = IF H2S > 0.01 THEN$$

$$\frac{kpim}{10} * rsop \quad (3.226)$$

$$ELSE kpim * rsop$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|-------------------------------------|--------|
| H2S | H ₂ S in the water above the sediment | g S m ⁻³ | S |
| kpim | Fraction of settled P to immobilisation | n.u. | C |
| rsop | Supply of organic P to sediment | g P m ⁻² d ⁻¹ | P |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.4.13 SNIM, Immobilised sediment N by denitrification & burial, g N m⁻²

$$\frac{dSNIM}{dt} = rsnim + rdenit \quad (3.227)$$

Where:

| Process | Comment | Unit |
|---------|--|-------------------------------------|
| rspim | Immobilisation of sediment N by burial | g N m ⁻² d ⁻¹ |
| rdenit | Denitrification of NO ₃ in sediment | g N m ⁻² d ⁻¹ |

rsnim: Immobilisation of sediment N by burial, g N m⁻²d⁻¹

Please see under state variable sediment organic N (SON), Section 3.4.5.

rdenit: Denitrification of NO₃ in sediment, g N m⁻²d⁻¹

Please see under state variable sediment NO₃ (SN03), Section 3.4.9.

3.4.14 SNIM, Immobilised sediment N by denitrification & burial, g N m⁻²

$$\frac{dSCIM}{dt} = minSOC + rscim \quad (3.228)$$

Where:

| Process | Comment | Unit |
|---------|------------------------------|-------------------------------------|
| minSOC | Mineralisation of SOC | g C m ⁻² d ⁻¹ |
| rscim | Burial of sediment organic C | g C m ⁻² d ⁻¹ |

minSOC: Mineralisation of SOC, g C m⁻²d⁻¹

Please see under state variable sediment organic C (SOC), Section 3.4.4.

rscim: Burial of sediment organic C, g C m⁻²d⁻¹

Please see under state variable sediment organic C (SOC), Section 3.4.4.

3.5 Help Processes

The help processes are divided into processes not included into differential equations (P1) and auxiliary processes (A). The distinction between is unimportant and rely on P1 processes in MIKE ECO Lab being able to be defined exclusively for bottom layer, surface layer and all layers in a 3D model. This distinction is not possible for auxiliary processes.

3.5.1 The P1 processes listed in alphabetic order

dkdox_no3: Change in NO₃ penetration rate sediment, analytical solution, m d⁻¹

NO₃ in the pore water of the sediment is caused by penetration of NO₃ from water above the sediment and by nitrification of NH₄ in the uppermost layer (KDO2) with DO in the pore water. The NO₃ concentration in this layer (0 to KDO2) has in the model the average concentration $SNO3m3$. From this layer NO₃ penetrates deeper into the anoxic sediment layer while being denitrified to N₂. Provided there is no outflow of NO₃ enriched ground water the NO₃ concentration at a sudden depth below the surface (KDOX) will be 0. Assuming a constant denitrification and under steady state condition the NO₃ concentration (NO3x) in the pore water at depth x below the surface layer with DO can be described with:

$$0 = -difno3 * \frac{d^2NO3x}{dx^2} + dnm3 \text{ Where } 0 < x < (KDOX_{\infty} - KDO2) \quad (3.229)$$

Which by integration becomes:

$$\frac{dNO3x}{dx} = \frac{dnm3}{difno3} * x + a \quad (3.230)$$

Where a is a constant, which by using the border condition ($dNO3x/dx=0$ at $x=(KDOX_{\infty} - KDO2)$) can be defined as:

$$a = -\frac{dnm3}{difno3} * (KDOX_{\infty} - KDO2) \Rightarrow \quad (3.231)$$

$$\frac{dNO3x}{dx} = \frac{dnm3}{difno3} * (x - (KDOX_{\infty} - KDO2))$$

Which by yet an integration gives:

$$NO3x = \frac{dnm3}{2*difno3} * x^2 - \frac{dnm3}{difno3} * (KDOX_{\infty} - KDO2) * x + b \quad (3.232)$$

Where b is a constant, which by using the border condition ($SNO3m3=0$ at $x=(KDOX_{\infty} - KDO2)$) can be defined as:

$$b = \frac{dnm3}{2*difno3} * (KDOX_{\infty} - KDO2)^2 \Rightarrow \quad (3.233)$$

$$NO3x = \frac{dnm3}{2*difno3} * x^2 - \frac{dnm3}{difno3} * (KDOX_{\infty} - KDO2) * x + \frac{dnm3}{2*difno3} * (KDOX_{\infty} - KDO2)^2$$

At depth $x=0$ in the anoxic zone (which is at depth $KDO2$ below sediment surface)
 $NO3x=SNO3m3$. =>

$$(KDOX_{\infty} - KDO2) = \sqrt{2 * difno3 * \frac{SNO3m3}{dnm3}} \Rightarrow$$

$$KDOX_{\infty} = \sqrt{2 * difno3 * \frac{SNO3m3}{dnm3}} + KDO2 \quad (3.234)$$

$KDOX_{\infty}$ in the above equation is the NO_3 penetration in the sediment under steady state condition. Assuming $KDO2_t - KDO2_{t+1}$ the change in the change in in $KDOX$ ($kdox_no3$) from time step t to time step $t+1$ can be defined as:

$$dkdox_{no3} = (KDOX_{\infty} - KDOX_t) * kkdox$$

$$dkdox_{no3} = \left(\sqrt{2 * difno3 * \frac{SNO3m3}{dnm3}} + KDO2_t - KDOX_t \right) * kkdox \quad (3.235)$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|---|--------------------|--------|
| difno3 | Vertical diffusion for NO_3 in sediment | m^2d^{-1} | C |
| dnm3 | denitrification in sediment, corrected for temperature | $g N m^{-3}d^{-1}$ | A |
| SNO3m3 | NO_3 in pore water surface sediment, layer (0- $kdo2$) | $g N m^{-3}$ | A |
| KDO2 | DO penetration into the sediment | m | S |
| KDO2 _t | DO penetration into the sediment , time step t | m | |
| KDOX | NO_3 penetration into the sediment, same as $KDOX_t$ | m | S |
| kkdox | Rate constant NO_3 penetration into sediment | d^{-1} | C |
| KDOX _t | NO_3 penetration into the sediment, time step t | m | |
| KDOX _∞ | NO_3 penetration into the sediment, steady state | m | |
| NO3x | NO_3 concentration in pore water at depth x | $g N m^{-3}$ | |
| x | Depth below zone with DO | m | |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fscb: Mineralisation of newly settled organic C, g C m⁻³d⁻¹

A fraction of newly settled particulate organic C (plankton and detritus) on the sediment surface is assumed to be mineralised at once. The fraction mineralised is dependent on the N:C ratio.

$$fscb = \frac{fsnb}{SEPN1 - BUOYN1 + SEPN2 + SEPN3 - BUOYN3 + SEDN} * \frac{depoC}{dz} \quad (3.236)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|-------------------------------------|--------|
| fsnb | Mineralisation of newly settled organic N | g N m ⁻³ d ⁻¹ | P1 |
| depoC | Deposition of particulate organic C | g C m ⁻² d ⁻¹ | P |
| SEPN1 | Sedimentation of flagellate N | g N m ⁻³ d ⁻¹ | P |
| SEPN2 | Sedimentation of diatom N | g N m ⁻³ d ⁻¹ | P |
| SEPN3 | Sedimentation of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| BUOYN1 | Flagellate upward movement N | g N m ⁻³ d ⁻¹ | P |
| BUOYN3 | Cyanobacteria upward movement N | g N m ⁻³ d ⁻¹ | P |
| SEDN | Deposition of detritus N | g N m ⁻³ d ⁻¹ | P |
| dz | Height of actual layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fsnb: Mineralisation of newly settled organic N, g N m⁻³d⁻¹

A fraction of newly settled particulate organic N (plankton, detritus and dead eelgrass) on the sediment surface is assumed to be mineralised at once. The fraction mineralised is dependent on the N:C ratio.

$$fsnb = krsn0 * \left(SEPN1 - BUOYN1 + SEPN2 + SEPN3 - BUOYN3 + SEDN - \frac{depoC * knim}{dz} \right) * tetn^{temp-20} \quad (3.237)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|-------------------------------------|-------------------------------------|--------|
| krsn0 | Fraction of deposited N mineralised | d ⁻¹ | C |
| depoC | Deposition of particulate organic C | g C m ⁻² d ⁻¹ | P |
| SEPN1 | Sedimentation of flagellate N | g N m ⁻³ d ⁻¹ | P |

| Name | Comment | Unit | Type*) |
|--------|---|-------------------------------------|--------|
| SEPN2 | Sedimentation of diatom N | g N m ⁻³ d ⁻¹ | P |
| SEPN3 | Sedimentation of cyanobacteria N | g N m ⁻³ d ⁻¹ | P |
| BUOYN1 | Flagellate upward movement N | g N m ⁻³ d ⁻¹ | P |
| BUOYN3 | Cyanobacteria upward movement N | g N m ⁻³ d ⁻¹ | P |
| SEDN | Deposition of detritus N | g N m ⁻³ d ⁻¹ | P |
| dz | Height of actual layer | m | F |
| knim | Sediment N:C ratio of immobile N | g N g C ⁻¹ | C |
| tetn | Θ value in Arrhenius temperature function | n.u. | C |
| temp | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fspb: Mineralisation of newly settled organic P, g P m⁻³d⁻¹

A fraction of newly settled particulate organic P (plankton and detritus) on the sediment surface is assumed to be mineralised at once. The fraction mineralised is not set to be dependent on the N:C ratio, because mineralised P as PO₄ can be adsorbed to resuspended fine sediment containing Fe⁺⁺⁺. The user should therefore consider this problem and set the constant *krsp0* accordingly, from model set up to model set up.

$$fspb = krsp0 * (SEPP1 - SUOYP1 + SEPP2 + SEPP3 - BUOYP3 + SEDP) * tetp^{temp-20} \quad (3.238)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|-------------------------------------|--------|
| krsp0 | Fraction of deposited N mineralised | d ⁻¹ | C |
| SEPP1 | Sedimentation of flagellate P | g P m ⁻³ d ⁻¹ | P |
| SEPP2 | Sedimentation of diatom P | g P m ⁻³ d ⁻¹ | P |
| SEPP3 | Sedimentation of cyanobacteria P | g P m ⁻³ d ⁻¹ | P |
| BUOYP1 | Flagellate upward movement P | g P m ⁻³ d ⁻¹ | P |
| BUOYP3 | Cyanobacteria upward movement P | g P m ⁻³ d ⁻¹ | P |
| SEDP | Deposition of detritus P | g P m ⁻³ d ⁻¹ | P |
| tetp | Θ value in Arrhenius temperature function | n.u. | C |
| temp | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

kdo2i: new steady state condition for KDO2, function of DO & respiration, analytical solution

From the sediment-water interface oxygen (DO) can penetrate into the sediment pore water by diffusion or actively being transported into the sediment by ventilation pumping and sediment mixing by the benthic fauna. Further microbenthic algae through photosynthesis can produce DO in the sediment-water interface. DO is consumed in the sediment by bacterial respiration and chemical oxidation of reduced substances (Fe^{++} , H_2S) resulting in the O_2 concentration becoming 0 (normally 0-2 cm) below the sediment surface. In the model this depth is defined as $KDO2$. Assuming the DO produced by the microbenthic algae is delivered to the water, the below differential equation can be set up assuming a steady state condition:

$$0 = -difo2 * \frac{d^2 O_2}{dy^2} + DOconsum \quad \text{Where } 0 < y < (KDO2_\infty) \quad (3.239)$$

Which by integration becomes:

$$\frac{d O_2}{dy} = \frac{DOconsum}{difo2} * y + a \quad (3.240)$$

Where a is a constant, which by using the border condition ($dO_2/dy=0$ at $y=KDO2_\infty$) can be defined as:

$$a = -\frac{DOconsum}{difo2} * KDO2_\infty \Rightarrow \quad (3.241)$$

$$\frac{d O_2}{dy} = \frac{DOconsum}{difo2} * y - \frac{DOconsum}{difo2} * KDO2_\infty$$

Which by yet an integration gives:

$$O_2 = \frac{DOconsum}{2 * difo2} * y^2 - \frac{DOconsum}{difo2} * KDO2_\infty * y + b \quad (3.242)$$

Where b is a constant, which by using the border condition ($O_2=0$ at $y=KDO2_\infty$) can be defined as:

$$b = \frac{DOconsum}{2 * difo2} * KDO2_\infty^2 \Rightarrow \quad (3.243)$$

$$O_2 = \frac{DOconsum}{2 * difo2} * y^2 - \frac{DOconsum}{difo2} * KDO2_\infty * y + \frac{DOconsum}{difo2} * KDO2_\infty^2$$

At the sediment surface $y=0$ the $O_2 = DO \Rightarrow$

$$KDO2_\infty = \sqrt{2 * difo2 * \frac{DO}{DOconsum}} \Rightarrow \quad (3.244)$$

$KDO2_\infty$ is identical to $ko2i$ in the model, however the DO consumption in the model is the sum of bacterial respiration ($reKDO2$), nitrification ($rsnit$) and a flux of reduced substances from the underlying sediment ($fsH2S$) to the layer with O_2 . All the mentioned DO consuming processes have the unit ($\text{g m}^{-2}\text{d}^{-1}$) and therefore have to be divided with the DO penetration from the previous time step t ($KDO2_t$). A conversion factor for $\text{O}_2:\text{N}$ of 4.57 $\text{g O}_2:\text{g NH}_4\text{-N}$ is used and a conversion factor for $\text{O}_2:\text{S}$ of 2 $\text{g O}_2:\text{g H}_2\text{S-S}$ is used.

The diffusion or rather transport of oxygen into the sediment is dependent of the activity of the benthic infauna. Their activity is linked to the DO concentration, at low DO (below 2 g m⁻³) the activity will decrease caused by increased mortality. The constant *difO2* is therefore multiplied by an oxygen function (1+sqdo).

$$kdo2i = \sqrt{2 * DO * \frac{difo2 * (1 + sqdo) * KDO2_t}{(rsnit * 4.57 + reKDO2 + fsh2s * 2)}} \quad (3.245)$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|---|--|--------|
| difo2 | Vertical diffusion for O ₂ in sediment, low fauna activity | m ² d ⁻¹ | C |
| DOconsum | Sediment O ₂ consumption, layer (0-KDO2) | g O ₂ m ⁻³ d ⁻¹ | |
| y | Depth below sediment surface | m | |
| KDO2 _∞ | DO penetration into the sediment, steady state= kdo2i | m | |
| KDO2 _t | DO penetration into the sediment time step t= KDO2 | m | |
| KDO2 | DO penetration into the sediment | m | S |
| sqdo | DO dependend auxiliary | n.u. | A |
| rsnit | Nitrification in sediment layer (0-KDO2) | g N m ⁻² d ⁻¹ | P |
| reKDO2 | DO consumption by bacteria layer (0-KDO2) | g O ₂ m ⁻² d ⁻¹ | P1 |
| fsh2s | Flux of SH ₂ S from reduced sediment to layer (0-KDO2) | g S m ⁻² d ⁻¹ | P |
| DO | O ₂ in water above sediment | g O ₂ m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

reKDO2: Sediment mineralisation of SOC by DO, in layer KDO2, g O₂ m⁻² d⁻¹

$$reKDO2 = \frac{KDO2}{kds} * minSOC * vo * sqdo \quad (3.246)$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|-------------------------------------|--------|
| KDO2 | DO penetration depth in sediment | m | S |
| kds | Depth of modelled sediment layer | m | C |
| minSOC | Mineralisation of organic in sediment | g C m ⁻² d ⁻¹ | P |
| vo | O ₂ : C ration production, respiration, mineralisation | g O ₂ g C ⁻¹ | C |
| sqdo | Oxygen function | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

3.5.2 Auxiliary (A) processes listed in alphabetic order

buoy1: N & P & light upward movement function, flagellate, n.u.

The upward movements of phytoflagellates are a function of the light regime and nutrient condition of the algae. If the internal N and P pools (N/C and P/C ratios in the algae) are small the function powNP1 will have a small but positive value resulting in an in sedbouNP1 being negative and the buoy1 becoming 0. At low powNP1 there will be no upward movements. At high powNP1 values (high N/C and P/C ratio in the algae) sedbouNP1 become positive and the algae will move upward provided the light doze is below a value of kiz3.

$$\begin{aligned} & buoy1 = IF \quad i \leq kiz3 \\ & \quad THEN \quad MAX(0, sedbouNP1) \\ & \quad ELSE \quad 0 \end{aligned} \quad (3.247)$$

Where:

$$sedbouNP1 = k2NP * \left(\frac{powNP1}{powNP1 + k3NP^{k1NP}} - 0.5 \right) \quad (3.248)$$

And:

$$powNP1 = MIN(myp1, myn1)^{k1NP} \quad (3.249)$$




Figure 3.3 Figure Flagellates upward movement is dependent of positive values of sedbuoy1 , which is dependent on a good N and P condition myn1 and myp1

Where:

| Name | Comment | Unit | Type*) |
|--------|--|---|--------|
| kiz3 | Light limit, buoyancy for PC1 & PC 3 | mol photon $\text{m}^{-2}\text{d}^{-1}$ | C |
| i | Photosynthetic Active Light (PAR) at top of layer | mol photon $\text{m}^{-2}\text{d}^{-1}$ | A |
| k1NP | Exponent, sedimentation & buoyancy, PC1&PC3 | n.u. | C |
| k2NP | Factor, sedimentation & buoyancy, PC1 & PC3 | n.u. | C |
| k3NP | Shift from sedimentation to buoyancy, PC1 & PC3 | n.u. | C |
| powNP1 | Power function limiting nutrient, Flagellate (PC1) | n.u. | A |
| myn1 | Nitrogen function flagellates | n.u. | A |
| myp1 | Phosphorous function flagellates | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

buoy3: N & P & light upward movement function, cyanobacteria, n.u.

The upward movement of cyanobacteria is a function of the light regime and nutrient condition of the bacteria. If the internal N and P pools (N/C and P/C ratios in the bacteria) are small the function powNP3 will have a small but positive value resulting in an in sedbouNP3 being negative and the buoy1 becoming 0. At low powNP3 there will be no upward movements. At high powNP3 values (high N/C and P/C ratio in the bacteria) sedbouNP3 become positive and the algae will move upward provided the light doze is below a value of kiz3.

$$buoy3 = IF \ i \leq kiz3 \ THEN \ MAX(0, sedbouNP3) \ ELSE \ 0 \tag{3.250}$$

Where:

$$sedbuoNP3 = k2NP * \left(\frac{powNP3}{powNP3 + k3NP^{k1NP}} - 0.5 \right) \tag{3.251}$$

And:

$$powNP3 = MIN(myp3, myn3)^{k1NP} \tag{3.252}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|--|--------|
| kiz3 | Light limit, buoyancy for PC1 & PC 3 | mol photon m ⁻² d ⁻¹ | C |
| i | Photosynthetic Active Light (PAR) at top of layer | mol photon m ⁻² d ⁻¹ | A |
| k1NP | Exponent, sedimentation& buoyancy, PC1&PC3 | n.u. | C |
| k2NP | Factor, sedimentation & buoyancy, PC1 & PC3 | n.u. | C |
| k3NP | Shift from sedimentation to buoyancy, PC1 & PC3 | n.u. | C |
| powNP3 | Power function limiting nutrient, cyanobacteria | n.u. | A |
| myn3 | Nitrogen function cyanobacteria | n.u. | A |
| myp3 | Phosphorous function cyanobacteria | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.




Figure 3.4 Cyanobacterial upward movement is dependent of positive values of sedbuoy3, which is dependent on a good N and P condition myn3 and myp3.

CSAIR: O₂ saturation in water, relative to PSU & temp., g O₂ m⁻³

A built in function in MIKE ECO Lab calculates the O₂ saturation relative to salinity and temperature. In this template the O₂ saturation defined by (Weiss 1970) is used:

$$CSAIR = \text{OXYGENSATURATION_WEISS}(S,T) \tag{3.253}$$

Or:

$$CSAIR = \frac{e^a}{0.69997} \tag{3.254}$$

Where:

$$a = -173.4292 + 249.6339 \frac{100}{T+273.15} + 143.2483 * \log\left(\frac{T+273.15}{100}\right) - 21.8493 * \left(\frac{T+273.15}{100}\right) + S * (-0.033096 + 0.014259 * \frac{T+273.15}{100} - 0.0017 * \left(\frac{T+273.15}{100}\right)^2) \tag{3.255}$$

In other templates the below equation is used.

$$CSAIR = \text{OXYGENSATURATION}(S,T) \tag{3.256}$$

Or:

$$CSAIR = 14.65 - 0.0841 \cdot S + T * (0.00256 \cdot S - 0.41022 + T * (0.007991 - 0.0000374 * S - 0.000077774 * T)) \tag{3.257}$$

Where:

| Name | Comment | Unit | Type*) |
|------|-------------|------|--------|
| S | Salinity | PSU | F |
| T | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

dndc: N:C ration, detritus, g N g C⁻¹

$$dndc = \frac{DN}{DC} \tag{3.258}$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------|---------------------|--------|
| DN | Detritus N | g N m ⁻³ | S |
| DC | Detritus C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

dnm3: denitrification in sediment, corrected for temperature

$$dnm3 = demax * tetn^{T-20} \tag{3.259}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|--|-------------------------------------|--------|
| Demax | Max. denitrification rate in sediment at 20 °C | g N m ⁻² d ⁻¹ | C |
| Tetn | Θ value in Arrhenius temperature function | n.u | C |
| T | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

doc_{monod}: UV radiation Monod relation for photo oxidation, n.u.

$$DOC_{monod} = \frac{i - doc_{ie}}{i - doc_{ie} + doc_{ik}} \quad (3.260)$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|--|--|--------|
| I | Solar radiation (PAR) in actual water column layer | μmol photon m ⁻² s ⁻¹ | A |
| doc _{ie} | min PAR light, CDOC photo oxidation | μmol photon m ⁻² s ⁻¹ | C |
| doc _{ik} | PAR half saturation photo oxidation of CDOC | μmol photon m ⁻² s ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

dpdc: P:C ration, detritus, g P g C⁻¹

$$dndc = \frac{DP}{DC} \quad (3.261)$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------|---------------------|--------|
| DP | Detritus P | g P m ⁻³ | S |
| DC | Detritus C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

dsidc: Si:C ration, detritus, g Si g C⁻¹

$$dSidc = \frac{DSi}{DC} \quad (3.262)$$

Where:

| Name | Comment | Unit | Type*) |
|------|-------------|----------------------|--------|
| DSi | Detritus Si | g Si m ⁻³ | S |
| DC | Detritus C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

DT_{day}: AD time step (and not HD time step!) in days, d step⁻¹

$$DT_{day} = \frac{DT}{86400} \tag{3.263}$$

Where:

| Name | Comment | Unit | Type*) |
|------|-------------------|-------------------------|--------|
| DT | Time step in sec. | Sec. step ⁻¹ | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

eta: Vertical light attenuation, m⁻¹

It is possible to calculate the vertical light attenuation based on light extinction constants or use the expression formulated by (Effer 1988; Kirk 2000) splitting the light attenuation into a light absorption and a scattering of light.

The surface area of the particles are important for the optical properties, as small particle have a larger surface area, they also have a higher vertical light extinction constant or a higher light scattering constant. In general the optical properties of particulate matter are proportional to the surface area of the particles in the water. In the equation for **eta1** and scattering **scatw** 3 size classes of inorganic matter (ss1-ss3) are defined in g m⁻³. The mass is not an ideal measure for inorganic matter in relation to the optical properties; therefore the mass of ss1-ss3 is related to a particle size with a diameter of 10 µm, by multiplication of a factor (10/diass_x). Small particles below 10 µm thereby will be assigned a higher light extinction constant, or absorption and scattering constants, whereas larger particles will be assigned smaller constants.

The shape of the particles may be anything from a ball to spherical cone; therefore the correction factor (10/diass_x) should only be regarded as guidelines to be used if no measured data exists.

In the present MIKE ECO Lab template resuspension is not included therefore the light extinction not included dynamically in the model often is put into the background light extinction (bla) and the light extinction from suspended matter is the light extinction from marine earth works. The user should in this case be aware that the background extinction varies in time and space especially in coastal waters.

$$eta = IF Keta > 0 THEN eta2 ELSE eta1 \tag{3.264}$$

Where:

$$eta1 = pla * CH + dla * DC + cla * CDOC + sla * 10 * \left(\frac{ss1}{diass1} + \frac{ss2}{diass2} + \frac{ss3}{diass3} \right) + bla \tag{3.265}$$

$$eta2 = \sqrt{absw^2 + 0.256 * absw * scatw}$$

And:

$$\text{absw} = \text{pla}_a * CH + \text{dla}_a * DC + \text{cla}_a * CDOC + \text{sla}_a * 10 * \left(\frac{\text{ss1}}{\text{diass1}} + \frac{\text{ss2}}{\text{diass2}} + \frac{\text{ss3}}{\text{diass3}} \right) + \text{bla}_a \quad (3.266)$$

$$\text{scatw} = \text{bkch} * CH^{\text{ekch}} + \text{bkss} * \left(\frac{10 * \text{ss1}}{\text{diass1}} + \frac{10 * \text{ss2}}{\text{diass2}} + \frac{10 * \text{ss3}}{\text{diass3}} \right)^{\text{ekss}}$$

The scatter of light (scatw) is defined as a power function of phytoplankton chlorophyll, (Morel A. 1980, Prieur L. & S. Sathyendranath 1981). The authors found a value range for bkch of 0.12-0.4 m²mg⁻¹ & ekch 0.63. The scatter by phytoplankton is dependent on cell size. The cell size tends to be smaller at low chlorophyll concentrations, where the plankton typically is nutrient limited (Yentsch C.S., D. A. Phinney 1989).

Lund-Hansen L.C. 2004 found chlorophyll in average to be responsible for 41% of the scattering in the nearby Århus Bay using a fixed specific chlorophyll scattering of 0.239 m²mg⁻¹ measured in New Zeland coastal waters (Pfanckuche F. 2002).

Please note that the mentioned scatter constants should be converted from m²mg⁻¹ to m²g⁻¹ before being used in this model.

Where:

| Name | Comment | Unit | Type*) |
|--------|--|--------------------------------|--------|
| Keta | Constant for choice of eta estimate | n.u. | C |
| eta1 | Function vertical light attenuation, extinction constants | m ⁻¹ | A |
| eta2 | Function vertical light attenuation, absorption & scattering | m ⁻¹ | A |
| pla | Chlorophyll light extinction constant | m ² g ⁻¹ | C |
| CH | Chlorophyll concentration | g m ⁻³ | S |
| dla | Detritus light extinction constant | m ² g ⁻¹ | C |
| DC | Detritus C | g m ⁻³ | S |
| cla | CDOC light extinction constant | m ² g ⁻¹ | C |
| CDOC | Coloured refractory DOC | g m ⁻³ | S |
| sla | Inorganic matter light extinction constant (Θ=10 μm) | m ² g ⁻¹ | C |
| ss1 | Inorganic matter, s e class 1 | g m ⁻³ | F |
| ss2 | Inorganic matter, size class 2 | g m ⁻³ | F |
| ss3 | Inorganic matter, size class 3 | g m ⁻³ | F |
| diass1 | Diameter of inorganic matter, size class 1 | μm | C |
| diass2 | Diameter of inorganic matter, size class 2 | μm | C |

| Name | Comment | Unit | Type*) |
|------------------|--|--------------------------------|--------|
| diass2 | Diameter of inorganic matter, size class 2 | µm | C |
| bla | Background light extinction | m ⁻¹ | C |
| absw | Light absorption in layer | m ⁻¹ | A |
| scatw | Light scattering in layer | m ⁻¹ | A |
| pla _a | Chlorophyll light absorption constant | m ² g ⁻¹ | C |
| dla _a | Detritus light absorption constant | m ² g ⁻¹ | C |
| cla _a | CDOC light absorption constant | m ² g ⁻¹ | C |
| sla _a | Inorganic matter light absorption constant (Θ=10 µm) | m ² g ⁻¹ | C |
| bla _a | Background light absorption | m ⁻¹ | C |
| bkch | Chlorophyll scattering constant | m ² g ⁻¹ | C |
| ekch | Chlorophyll scattering exponent | n.u. | C |
| bkss | Inorganic matter scattering constant | m ² g ⁻¹ | C |
| ekss | Inorganic matter scattering exponent | n.u. | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fiz: Light factor for flagellates and cyanobacteria (PC1 & PC3) sedimentation, n.u.

$$fiz = fiz1 * fiz2 \tag{3.267}$$

Where:

$$fiz1 = IF i > kiz1 THEN 3 ELSE 1 \tag{3.268}$$

$$fiz2 = IF i > kiz2 THEN 1 ELSE 0$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|--|--------|
| fiz1 | 1. Help factor for PC1 & PC3 sedimentation | n.u. | A |
| fiz2 | 2. Help factor for PC1 & PC3 sedimentation | n.u. | A |
| i | Light at top of actual water layer | mol photon m ⁻² d ⁻¹ | A |
| kiz1 | Light limit for 3 X sedimentation rate of PC1 & PC3 | mol photon m ⁻² d ⁻¹ | C |
| kiz2 | Light limit for 1 X sedimentation rate of PC1 & PC3 | mol photon m ⁻² d ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fiz1: 1. Help factor for PC1 & PC3 sedimentation, n.u.

See under auxiliary fiz, Equation (3.18) and Equation (3.29).

fiz2: 2. Help factor for PC1 & PC3 sedimentation, n.u.

See under auxiliary fiz, Equation (3.18) and Equation (3.29).

fn3a: Denitrification, DO dependency in water column, n.u.

$$fn3a = \frac{ksb}{ksb + DO} \tag{3.269}$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|----------------------------------|--------|
| ksb | Denitrification Half saturation conc. DO | g O ₂ m ⁻³ | C |
| DO | Oxygen concentration | g O ₂ m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fp3sal: Function for cyanobacteria dependency (death & growth) of salinity, n.u.

$$fp3sa = IF S < kp3opti THEN 1 ELSE e^{-kp3sal1*(S-kp3sal2)} \tag{3.270}$$

Where:

| Name | Comment | Unit | Type*) |
|---------|--|-------------------|--------|
| kp3opti | Highest salinity for optimum cyanobacteria growth | PSU | C |
| kp3sal1 | Cyanobacteria growth salinity dependency coefficient | PSU ⁻¹ | C |
| kp3sal2 | Cyanobacteria growth salinity dependency constant | PSU | C |
| S | Salinity | PSU | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

fsa: Salinity function for reduction of SO₄ to H₂S, n.u.

$$fsa = IF S > ksa THEN 1 ELSE 0 \tag{3.271}$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|------|--------|
| ksa | Minimum salinity for SO ₄ reduction | PSU | C |
| S | Salinity | PSU | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

i: Solar radiation (PAR) in actual water column layer (j), mol photon m⁻²d⁻¹

If the surface water temperature >0.2 °C the water is assumed ice free and i₀ will be the light (PAR) reaching the surface of the water.

If the temperature is below 0.2 °C ice is assume on the water and only 10% of i₀ is assumed to penetrate the ice cover.

In a more general form the light (PAR) distribution in the different water layer can be expressed as:

$$i = i_0 * e^{\sum_j^0 - eta_{0-j} * dz_{0-j}} \tag{3.272}$$

MIKE ECO Lab has an builtin function ([LAMBERT_BEER_1](#)) calculate the light (PAR) at the top of each water layer.

The average light (PAR) in a water layer can be expressed as:

$$i = \frac{1 - e^{-eta * dz}}{eta * dz} * LAMBERT_BEER_1(i_0, dz, eta) \tag{3.273}$$

Where:

| Name | Comment | Unit | Type*) |
|--------------------|---|--|--------|
| i ₀ | Solar (PAR) radiation at water surface | mol photon m ⁻² d ⁻¹ | F |
| eta _{0-j} | Light attenuation (Kd) in layers 0 to j | m ⁻¹ | |
| dz _{0-j} | Height of layer 0 to j | m | |
| eta | Light attenuation (Kd) in actual layer | m ⁻¹ | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

ik1: Temperature corrected Light saturation for flagellates, mol photon m⁻²d⁻¹

$$ik1 = \text{alfa1} * \text{teti}^{T-20} \quad (3.274)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|--|--------|
| alfa1 | Light saturation at 20 °C for flagellates | mol photon m ⁻² d ⁻¹ | C |
| teti | Θ value Arrhenius expression | n.u | C |
| T | temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

ik2: Temperature corrected Light saturation for diatoms, mol photon m⁻²d⁻¹

$$ik2 = \text{alfa2} * \text{teti}^{T-20} \quad (3.275)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---------------------------------------|--|--------|
| Alfa2 | Light saturation at 20 °C for diatoms | mol photon m ⁻² d ⁻¹ | C |
| teti | Θ value Arrhenius expression | n.u | C |
| T | temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

ik3: Temperature corrected Light saturation for cyanobacteris, mol photon m⁻²d⁻¹

$$ik3 = \text{alfa3} * \text{teti}^{T-20} \quad (3.276)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|--|--------|
| Alfa3 | Light saturation at 20 °C for cyanobacteria | mol photon m ⁻² d ⁻¹ | C |
| teti | Θ value Arrhenius expression | n.u | C |
| T | temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

ksd: Sedimentation rate detritus, d⁻¹

The sedimentation of a particle is not allowed to pass through several water layers in one time step (DT_{day} time step in d). The sedimentation in one times step is therefore restricted to a maximum of dz.

This restriction may led to an underestimation of sedimentation in 3-D set ups with a fine vertical resolution using small dz. However by not imposing the below restriction in sedimentation will potentially generate mass balance errors.

$$ksd = IF dz \leq sevd * DT_{day} THEN \frac{dz}{DT_{day}} ELSE sevd \tag{3.277}$$

Where:

| Name | Comment | Unit | Type*) |
|-------------------|------------------------------|----------------------|--------|
| dz | Height of actual water layer | m | F |
| DT _{day} | AD time step | d step ⁻¹ | C |
| sevd | Sedimentation rate | m d ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mgpc: Temperature, O₂ & food corrected max. grazing rate by zooplankton, d⁻¹

$$mgpc = kgrb * tetz^{T-20} * \frac{sqdo}{1 + e^{kgrm - kgrs * (kedib1 * PC1 + kedib2 * PC2 + kedib3 * PC3)}} \tag{3.278}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|-----------------|--------|
| kgrb | Max. specific grazing rate, zooplankton | d ⁻¹ | C |
| tetz | Θ in Arrhenius temp. relation of zooplankton grazing | n.u. | C |
| T | Temperature | °C | F |
| sqdo | DO function | n.u. | A |
| kgrm | Zooplankton 0. order dependency of grazing on plankton | n.u. | C |
| kgrs | Zooplankton 1. order dependency of grazing on plankton | n.u. | C |
| kedib1 | Edible fraction of Flagellate | n.u. | C |
| kedib2 | Edible fraction of Flagellate | n.u. | C |

| Name | Comment | Unit | Type*) |
|--------|-------------------------------|---------------------|--------|
| kedib3 | Edible fraction of Flagellate | n.u. | C |
| PC1 | Flagellate C | g C m ⁻³ | S |
| PC2 | Diatom C | g C m ⁻³ | S |
| PC3 | Cyanobacteria C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mnl1: Nutrient dependent death factor, flagellate, n.u.

The death factor is 1 with flagellate having high internal N:C and P:C ratios, but up to 5 in a nutrient stressed condition (low N:C and P:C ratios).

$$mnl1 = MIN\left(\frac{\left(\frac{1}{myn1} + \frac{1}{myp1}\right)}{2}, 5\right) \quad (3.279)$$

Where:

| Name | Comment | Unit | Type*) |
|------|---------------------------------|------|--------|
| myn1 | Nitrogen function flagellate | n.u | A |
| myp1 | Phosphorous function flagellate | n.u | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mnl2: Nutrient dependent death factor, diatoms, n.u.

The death factor is 1 with diatoms having high internal N:C, P:C and Si:C ratios, but up to 5 in a nutrient stressed condition (low N:C, P:C or Si:C ratios).

$$mnl2 = MIN\left(\frac{\left(\frac{1}{myn2} + \frac{1}{myp2} + \frac{1}{mys2}\right)}{3}, 5\right) \quad (3.280)$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------------|------|--------|
| myn2 | Nitrogen function diatoms | n.u | A |
| myp2 | Phosphorous function diatoms | n.u | A |
| mys2 | Si function, diatoms | n.u | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mnl3: Nutrient dependent death factor, cyanobacteria, n.u.

The death factor is 1 with cyanobacteria having high internal N:C and P:C ratios, but up to 5 in a nutrient stressed condition (low N:C and P:C ratios).

$$mnl3 = MIN\left(\frac{\left(\frac{1}{myn3} + \frac{1}{myp3}\right)}{2}, 5\right) \tag{3.281}$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------------------|------|--------|
| myn3 | Nitrogen function cyanobacteria | n.u | A |
| myp3 | Phosphorous function cyanobacteria | n.u | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mntp1: N, P & temperature corrected max. net growth rate, flagellates d⁻¹

$$mntp1 = myte1 * \frac{2}{\left(\frac{1}{myn1} + \frac{1}{myp1}\right)} \tag{3.282}$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|------|--------|
| myte1 | Specific growth ,temperature regulated, flagellates | n.u. | A |
| myn1 | Nitrogen function flagellates | n.u. | A |
| myp1 | Phosphorous function flagellates | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mntp2: N, P & temperature corrected max. net growth rate, diatoms d⁻¹

$$mntp2 = myte2 * \frac{3}{\left(\frac{1}{myn2} + \frac{1}{myp2} + \frac{1}{mys2}\right)} \quad (3.283)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|------|--------|
| myte2 | Specific growth ,temperature regulated, diatoms | n.u. | A |
| myn2 | Nitrogen function diatoms | n.u. | A |
| myp2 | Phosphorous function diatoms | n.u. | A |
| mys2 | Silicate function diatoms | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mntp3: N, P & temperature corrected max. net growth rate, cyanobacteria d⁻¹

$$mntp3 = myte3 * \frac{2}{\left(\frac{1}{myn3} + \frac{1}{myp3}\right)} \quad (3.284)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|------|--------|
| myte3 | Specific growth ,temperature regulated, cyanobacteria | n.u. | A |
| myn3 | Nitrogen function cyanobacteria | n.u. | A |
| myp3 | Phosphorous function cyanobacteria | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mssp1: Sedimentation rate flagellate phytoplankton, d⁻¹

The sedimentation of a particle is not allowed to pass through several water layers in one time step (DT_{day} time step in d). The sedimentation in one times step is therefore restricted to a maximum of dz.

$$mssp1 = IF dz \leq seve1 * DT_{day} THEN \frac{dz}{DT_{day}} ELSE seve1 \quad (3.285)$$

This restriction may lead to an underestimation of sedimentation in 3D set-ups with a fine vertical resolution using small dz. However, by not imposing the above restriction in sedimentation will potentially generate mass balance errors.

Where:

| Name | Comment | Unit | Type*) |
|-------------------|---|-------------------|--------|
| seve1 | Sedimentation rate flagellate | m d ⁻¹ | C |
| DT _{day} | AD time step (and not HD time step!) in days | d | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

m_{spc2}: Sedimentation rate diatom phytoplankton, d⁻¹

The sedimentation of a particle is not allowed to pass through several water layers in one time step (DT_{day} time step in d). The sedimentation in one times step is therefore restricted to a maximum of dz.

$$m_{spc2} = IF dz \leq seve2 * DT_{day} THEN \frac{dz}{DT_{day}} ELSE seve2 \quad (3.286)$$

This restriction may lead to an underestimation of sedimentation in 3D set-ups with a fine vertical resolution using small dz. However, by not imposing the above restriction in sedimentation will potentially generate mass balance errors.

Where:

| Name | Comment | Unit | Type*) |
|-------------------|---|-------------------|--------|
| seve2 | Sedimentation rate diatoms | m d ⁻¹ | C |
| DT _{day} | AD time step (and not HD time step!) in days | d | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

m_{spc3}: Sedimentation rate cyanobacteria, d⁻¹

The sedimentation of a particle is not allowed to pass through several water layers in one time step (DT_{day} time step in d). The sedimentation in one times step is therefore restricted to a maximum of dz.

$$m_{spc3} = IF dz \leq seve3 * DT_{day} THEN \frac{dz}{DT_{day}} ELSE seve3 \quad (3.287)$$

This restriction may lead to an underestimation of sedimentation in 3D set-ups with a fine vertical resolution using small dz. However, by not imposing the above restriction in sedimentation will potentially generate mass balance errors.

Where:

| Name | Comment | Unit | Type*) |
|-------------------|---|-------------------|--------|
| seve3 | Sedimentation rate cyanobacteria | m d ⁻¹ | C |
| DT _{day} | AD time step (and not HD time step!) in days | d | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myfi1: Light function Flagellate, n.u.

Equation (3.288) is an analytical solution of the integrated light available for production of flagellates in the water column.

Where:

$$myfi1 = (zk1 + \frac{i}{ik1 * eta} * (e^{-eta*zk1} - e^{-eta*dz}))/dz \quad (3.288)$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|--|--------|
| zk1 | Light availability flagellate production | m | A |
| ik1 | Light saturation flagellate temperature corrected | mol photon m ⁻² d ⁻¹ | A |
| eta | Vertical light attenuation | m ⁻¹ | A |
| i | Photosynthetic Active Light (PAR) of layer | mol photon m ⁻² d ⁻¹ | A |
| dz | Height of actual layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myfi2: Light function diatom, n.u.

Equation (3.289) is an analytical solution of the integrated light available for production of flagellates in the water column.

$$myfi2 = (zk2 + \frac{i}{ik2 * eta} * (e^{-eta*zk2} - e^{-eta*dz}))/dz \quad (3.289)$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|---|--------|
| zk2 | Light availability diatom production | m | A |
| ik2 | Light saturation diatome temperature corrected | mol photon m ⁻² d ⁻¹ | A |
| eta | Vertical light attenuation | m ⁻¹ | A |
| i | Photosynthetic Active Light (PAR) of layer | mol photon m ⁻² d ⁻¹ | A |
| dz | Height of actual layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myfi3: Light function cyanobacteria, n.u.

Equation (3.290) is an analytical solution of the integrated light available for production of flagellates in the water column.

$$myfi3 = (zk3 + \frac{i}{ik3 * eta} * (e^{-eta*zk3} - e^{-eta*dz}))/dz \tag{3.290}$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|---|--------|
| zk3 | Light availability cyanobacteria production | m | A |
| ik3 | Light saturation cyanobacteria temp. corrected | mol photon m ⁻² d ⁻¹ | A |
| eta | Vertical light attenuation | m ⁻¹ | A |
| i | Photosynthetic Active Light (PAR) of layer | mol photon m ⁻² d ⁻¹ | A |
| dz | Height of actual layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myn1: Nitrogen function flagellates, n.u.

$$myn1 = \frac{PN1/PC1 - pnmi}{(pnma - pnmi)} \quad (3.291)$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------------------|-----------------------|--------|
| PC1 | Flagellate phytoplankton C | g C m ⁻³ | S |
| PN1 | Flagellate phytoplankton N | g N m ⁻³ | S |
| pnmi | Minimum N:C ratio in phytoplankton | g N g C ⁻¹ | C |
| pnma | Maximum N:C ratio in phytoplankton | g N g C ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myn2: Nitrogen function Diatoms, n.u.

$$myn2 = \frac{\frac{PN2}{PC2} - psmi * pnsi}{(pnma - psmi * pnsi)} \quad (3.292)$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------------------|------------------------|--------|
| PC2 | Diatom phytoplankton C | g C m ⁻³ | S |
| PN2 | Diatom phytoplankton N | g N m ⁻³ | S |
| psmi | Minimum Si:C ratio in diatoms | g Si g C ⁻¹ | C |
| pnsi | Minimum N:Si ratio in diatoms | g N g Si ⁻¹ | C |
| pnma | Maximum N:C ratio in phytoplankton | g N g C ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myn3: Nitrogen function cyanobacteria, n.u.

$$myn3 = \frac{PN3/PC3 - pnmi}{(pnma - pnmi)} \quad (3.293)$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|-----------------------|--------|
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| PN3 | Cyanobacteria N | g N m ⁻³ | S |
| pnmi | Minimum N:C ratio in phytoplankton / cyanobacteria | g N g C ⁻¹ | C |
| pnma | Maximum N:C ratio in phytoplankton / cyanobacteria | g N g C ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myp1: Phosphorous function flagellates, n.u.

$$myp1 = \frac{(kc + ppma - ppmi) * (\frac{PP1}{PC1} - ppmi)}{(ppma - ppmi) * (kc + \frac{PP1}{PC1} - ppmi)} \tag{3.294}$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|-----------------------|--------|
| PC1 | Flagellate phytoplankton C | g C m ⁻³ | S |
| PP1 | Flagellate phytoplankton P | g P m ⁻³ | S |
| ppmi | Minimum P:C ratio in phytoplankton | g P g C ⁻¹ | C |
| ppma | Maximum P:C ratio in phytoplankton | g P g C ⁻¹ | C |
| kc | Half saturation concentration for phytoplankton P | g P g C ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myp2: Phosphorous function diatoms, n.u.

$$myp2 = \frac{(kc + ppma - ppmi) * (\frac{PP2}{PC2} - ppmi)}{(ppma - ppmi) * (kc + \frac{PP2}{PC2} - ppmi)} \tag{3.295}$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------------------|-----------------------|--------|
| PC2 | Flagellate phytoplankton C | g C m ⁻³ | S |
| PP2 | Flagellate phytoplankton P | g P m ⁻³ | S |
| ppmi | Minimum P:C ratio in phytoplankton | g P g C ⁻¹ | C |

| Name | Comment | Unit | Type*) |
|------|---|-----------------------|--------|
| ppma | Maximum P:C ratio in phytoplankton | g P g C ⁻¹ | C |
| kc | Half saturation concentration for phytoplankton P | g P g C ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mys3: Phosphorous function cyanobacteria, n.u.

$$mys3 = \frac{\left(\frac{kc}{ppma} * p3pma + p3pma - p3pmi\right) * \left(\frac{PP3}{PC3} - p3pmi\right)}{\left(p3pma - p3pmi\right) * \left(\frac{kc}{ppma} * p3pma + \frac{PP3}{PC3} - p3pmi\right)} \quad (3.296)$$

Where:

| Name | Comment | Unit | Type*) |
|-------|---|-----------------------|--------|
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| PP3 | Cyanobacteria P | g P m ⁻³ | S |
| ppmi | Minimum P:C ratio in phytoplankton | g P g C ⁻¹ | C |
| ppma | Maximum P:C ratio in phytoplankton | g P g C ⁻¹ | C |
| kc | Half saturation conc. for phytoplankton | g P g C ⁻¹ | C |
| p3pma | Maximum P:C ratio in cyanobacteria | g P g C ⁻¹ | C |
| p3pmi | Minimum P:C ratio in cyanobacteria | g P g C ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

mys2: Si function, diatoms, n.u

$$mys2 = \frac{\frac{PSi2}{PC2} - psmi}{(psma + psmi)} \quad (3.297)$$

Where:

| Name | Comment | Unit | Type*) |
|------|-------------------------------|------------------------|--------|
| PC2 | Diatom phytoplankton C | g C m ⁻³ | S |
| PSi2 | Diatom phytoplankton Si | g Si m ⁻³ | S |
| psmi | Minimum Si:C ratio in diatoms | g Si g C ⁻¹ | C |
| psma | Maximum Si:C ratio in diatoms | g Si g C ⁻¹ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myte1: Flagellate specific temperature corrected growth, d⁻¹

$$myte1 = mym1 * tet1^{T-20} \tag{3.298}$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|-----------------|--------|
| mym1 | Max. specific net growth at 20 °C, flagellates | d ⁻¹ | C |
| tet1 | Θ value in Arrhenius relation, flagellate temp. relation | n.u. | C |
| T | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myte2: Diatom specific temperature corrected growth, d⁻¹

Smoot_T is a function that makes a smoothing or rolling average of the daily surface insolation (PAR) over a number of days (T2_{days}) defined by the user, see auxiliary Smoot_T, Equation (3.318). This smoothing or rolling average is used to adjust the reference temperature of the max. specific growth rate of the diatoms.

The seasonal variation of temperature in the water follow the seasonal variation of the light (PAR) with a delay (1 month) depending of the amount of water (depth) to be heated up. In spring the temperature will be low compared to the daily PAR doze whereas in fall the temperature will be high compared to the daily PAR doze. The diatom reference temperature therefor has to change over the season.

In spring the diatoms blooms at low temperatures and disappear when the silicate is used up. During summer some silicate will be available however the diatom community has changed and another higher reference temperature is needed. In fall a secondary diatom bloom is sometimes seen after the erosion of the pycnocline. The diatom community is again adapted to lower temperatures and decreasing PAR.

Introducing Smoot_T is an attempt to make a seasonal adjustment of the specific growth with the water temperature and light as forcing.

$$myte2 = mym2 * tet2^{T-6-Smoot_T} \quad (3.299)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|--|-----------------|--------|
| mym2 | Max. specific net growth at 6-10 °C, Diatoms | d ⁻¹ | C |
| tet2 | Θ value in Arrhenius relation, diatom temp. relation | n.u. | C |
| Smoot_T | Correction of reference temp. for diatoms | °C | A |
| T | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

myte3: Cyanobacteria specific temperature corrected growth, d⁻¹

$$myte3 = mym3 * tet3^{T-20} \quad (3.300)$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|-----------------|--------|
| mym3 | Max. specific net growth at 20 °C, cyanobacteria | d ⁻¹ | C |
| tet3 | Θ value in Arrhenius relation, cyanobacteria temp. relation | n.u. | C |
| T | Temperature | °C | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

nfix1: Function for N fixation (1 if PSU≤12 else 0), n.u.

$$nfix1 = IF S \leq 12 THEN 1 ELSE 0 \quad (3.301)$$

Where:

| Name | Comment | Unit | Type*) |
|------|----------|------|--------|
| S | Salinity | PSU | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

nfix2: Function for N fixation (1 if 0≤PSU≤10 else 0-1), n.u.

$$nfix2 = IF S \geq 12 THEN (1 - \frac{S - 10}{12 - 10}) ELSE 1 \tag{3.302}$$

Where:

| Name | Comment | Unit | Type*) |
|------|----------|------|--------|
| S | Salinity | PSU | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

pn1pc1: N:C ration flagellates g N g C⁻¹

$$pn1pc1 = \frac{PN1}{PC1} \tag{3.303}$$

Where:

| Name | Comment | Unit | Type*) |
|------|----------------------------|---------------------|--------|
| PN1 | Flagellate phytoplankton N | g N m ⁻³ | S |
| PC1 | Flagellate phytoplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

pn2pc2: N:C ration diatoms g N g C⁻¹

$$pn2pc2 = \frac{PN2}{PC2} \tag{3.304}$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------|---------------------|--------|
| PN2 | Diatom phytoplankton N | g N m ⁻³ | S |
| PC2 | Diatom phytoplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

pn3pc3: N:C ration cyanobacteria g N g C⁻¹

$$pn3pc3 = \frac{PN3}{PC3} \quad (3.305)$$

Where:

| Name | Comment | Unit | Type*) |
|------|-----------------|---------------------|--------|
| PN3 | Cyanobacteria N | g N m ⁻³ | S |
| PC3 | Cyanobacteria C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

powNP1: Power function for limiting nutrient, Flagellate (PC1), n.u.

See under auxiliary sed1, Equation (3.18).

powNP3: Power function for limiting nutrient, cyanobacteria (PC3), n.u.

See under auxiliary sed3, Equation (3.29).

pp1pc1: P:C ration flagellates g P g C⁻¹

$$pp1pc1 = \frac{PP1}{PC1} \quad (3.306)$$

Where:

| Name | Comment | Unit | Type*) |
|------|----------------------------|---------------------|--------|
| PP1 | Flagellate phytoplankton P | g P m ⁻³ | S |
| PC1 | Flagellate phytoplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

pp2pc2: P:C ration diatoms g P g C⁻¹

$$pp2pc2 = \frac{PP2}{PC2} \quad (3.307)$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------|---------------------|--------|
| PP2 | Diatom phytoplankton P | g P m ⁻³ | S |
| PC2 | Diatom phytoplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

pp3pc3: P:C ration cyanobacteria g P g C⁻¹

$$pp3pc3 = \frac{PP3}{PC3} \tag{3.308}$$

Where:

| Name | Comment | Unit | Type*) |
|------|-----------------|---------------------|--------|
| PP3 | Cyanobacteria P | g P m ⁻³ | S |
| PC3 | Cyanobacteria C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

psi2pc2: Si:C ration diatoms g Si g C⁻¹

$$psi2pc2 = \frac{PSi2}{PC2} \tag{3.309}$$

Where:

| Name | Comment | Unit | Type*) |
|------|------------------------|----------------------|--------|
| PSi2 | Diatom Si | g Si m ⁻³ | S |
| PC2 | Diatom phytoplankton C | g C m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

rd: Relative daylength, f(month, day,latitude), n.u.

A built in function in MIKE ECO Lab that returns a value for relative day length. The value equals 1 at equinox (when day and night have same length).

$$rd = RELATIVE_DAYLENGTH(month, day, latitude) \tag{3.310}$$

sedbouNP1: sedimentation & buoyance N&P function, Flagellate, n.u

Please see auxiliary sed1, Equation (3.18), or buoy1, Equation (3.19)

sedbouNP3: sedimentation & buoyance N&P function, cyanobacteria, n.u

Please see auxiliary sed3, Equation (3.29), or buoy3, Equation (3.30).

sed1: N&P sedimentation function, Flagellate, n.u.

The sedimentation (downward movement) of the algae is increased by low internal N/C and P/C ratios of the algae. powNP1 will be positive but small and sedbouNP1 becomes negative resulting in a positive value of sed1, see Figure 3.3, under auxiliary buoy1.

$$sed1 = MAX(0, -sedbouNP1) \tag{3.311}$$

Where:

$$sedbouNP1 = k2NP * \left(\frac{powNP1}{powNP1 + k3NP^{k1NP}} - 0.5 \right) \tag{3.312}$$

And:

$$powNP1 = MIN(myp1, myn1)^{k1NP} \tag{3.313}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|--|------|--------|
| k1NP | Exponent, sedimentation & buoyancy, PC1&PC3 | n.u. | C |
| K2NP | Factor for sedimentation & buoyancy, PC1 & PC3 | n.u. | C |
| k3NP | Shift from sedimentation to buoyancy, PC1 & PC3 | n.u. | C |
| powNP1 | Power function for limiting nutrient, Flagellate (PC1) | n.u. | A |
| myn1 | Nitrogen function flagellates | n.u. | A |
| myp1 | Phosphorous function flagellates | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

Sed3: N&P sedimentation function, cyanobacteria, n.u.

The sedimentation (downward movement) of the bacteria is increased by low internal N/C and P/C ratios of the bacteria. powNP3 will be positive but small and sedbouNP3 becomes negative resulting in a positive value of sed3, see Figure 3.4 under auxiliary bou3.

$$sed3 = MAX(0, -sedbuoNP3) \tag{3.314}$$

Where:

$$sedbuoNP3 = k2NP * (\frac{powNP3}{powNP3 + k3NP^{k1NP}} - 0.5) \tag{3.315}$$

And:

$$powNP3 = MIN(myp3, myn3)^{k1NP} \tag{3.316}$$

Where:

| Name | Comment | Unit | Type*) |
|--------|---|------|--------|
| k1NP | Exponent, sedimentation & buoyancy, PC1&PC3 | n.u. | C |
| K2NP | Factor for sedimentation & buoyancy, PC1 & PC3 | n.u. | C |
| k3NP | Shift from sedimentation to buoyancy, PC1 & PC3 | n.u. | C |
| powNP3 | Power function for limiting nutrient, cyanobacteria (PC3) | n.u. | A |
| myn3 | Nitrogen function cyanobacteria | n.u. | A |
| myp3 | Phosphorous function cyanobacteria | n.u. | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

SIPm3: PO₄-P in pore water, g P m⁻³

$$SIPm3 = \frac{SIP}{(1 - dm) * vf * kds} \tag{3.317}$$

Where:

| Name | Comment | Unit | Type*) |
|------|----------------------------------|------------------------|--------|
| SIP | Sediment PO ₄ -P pool | g P m ⁻² | S |
| dm | Sediment dry matter | g DM gWW ⁻¹ | C |
| vf | Sediment bulk density | g ww cm ⁻³ | C |
| Kds | Depth of modelled sediment layer | m | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

Smoot_T: Correction of reference temperature for diatoms, °C

Smoot_T is a function that makes a smoothing or rolling average of the daily insolation (PAR) on surface over a number of days (T_{2days}) defined by the user. This smoothing or rolling average is used to adjust the reference temperature of the max. specific growth rate of the diatoms, see auxiliary *myte2*, Equation (3.299).

The seasonal variation of temperature in the water follow the seasonal variation of the light (PAR) with a delay (1 month) depending of the amount of water (depth) to be heated up. In spring the temperature will be low compared to the daily PAR doze whereas in fall the temperature will be high compared to the daily PAR doze. The diatom reference temperature therefor has to change over the season.

In spring the diatoms blooms at low temperatures and disappear when the silicate is used up. During summer some silicate will be available however the diatom community has changed and another higher reference temperature is needed. In fall a secondary diatom bloom is sometimes seen after the erosion of the pycnocline. The diatom community is again adapted to lower temperatures and decreasing PAR.

Introducing Smoot_T is an attempt to make a seasonal adjustment of the specific growth with the water temperature and light as forcing.

One of two builtin functions can be used SMOOTHING_AVERAGE or MOVING_AVERAGE can be used, see (MIKE by DHI 2011b). The latter function demands more memory and increases the CPU time slightly.

$$Smoot_T = \frac{difT2}{maxI_0} * SMOOTHING_AVERAGE(i_0, \frac{DT_{day}}{T2_{days}}) \quad (3.318)$$

Where:

| Name | Comment | Unit | Type*) |
|--------------------|---|---|--------|
| difT2 | Max variation in reference temp., diatom production | °C | C |
| maxI ₀ | Max average monthly i ₀ of year (July or Jan.) | mol photon m ⁻² d ⁻¹ | C |
| i ₀ | Light (PAR) at surface | mol photon m ⁻² d ⁻¹ | F |
| DT _{day} | AD time step in days, (normally between 5 min. to 1 h) | d | F |
| T2 _{days} | No. of days in smoothing or rolling average function | d | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

SNHm3: NH₄-N in sediment pore water, g N m⁻³

$$SNHm3 = \frac{SNH}{(1 - dm) * vf * kds} \quad (3.319)$$

Where:

| Name | Comment | Unit | Type*) |
|------|----------------------------------|------------------------|--------|
| SNH | Sediment NH ₄ -N pool | g N m ⁻² | S |
| dm | Sediment dry matter | g DM gWW ⁻¹ | C |
| vf | Sediment bulk density | g ww cm ⁻³ | C |
| Kds | Depth of modelled sediment layer | m | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

SNO3m3: NO₃-N in pore water of surface sediment layer (0-KDO₂) , g N m⁻³

$$SNO3m3 = \frac{SNO3}{(1 - dm) * vf * KDO2} \quad (3.320)$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|------------------------|--------|
| SNO3 | Sediment NO ₃ -N pool | g N m ⁻² | S |
| dm | Sediment dry matter | g DM gWW ⁻¹ | C |
| vf | Sediment bulk density | g ww cm ⁻³ | C |
| KDO2 | Depth of O ₂ penetration in sediment | m | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

sqdo: Oxygen function, n.u.

$$sqdo = \frac{DO^{ndo3}}{DO^{ndo3} + mdo3} \quad (3.321)$$

Where:

| Name | Comment | Unit | Type*) |
|------|-----------------------------|----------------------------------|--------|
| DO | Oxygen | g O ₂ m ⁻³ | S |
| ndo3 | Exponent for DO in sqdo | n.u. | C |
| mdo3 | Half-saturation constant DO | g O ₂ m ⁻³ | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

zk1: Light penetration (availability) in actual layer for flagellate production, m

$$zk1 = MIN(dz, MAX(0, \frac{\ln(i) - \ln(ik1)}{\eta})) \quad (3.322)$$

Where:

| Name | Comment | Unit | Type*) |
|------|--|--|--------|
| i | Light (PAR) in actual layer | mol photon m ⁻² d ⁻¹ | A |
| ik1 | Light saturation temp. corrected, Flagellate | mol photon m ⁻² d ⁻¹ | A |
| eta | Vertical light extinction in layer | m ⁻¹ | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

zk2: Light penetration (availability) in actual layer for diatom production, m

$$zk1 = MIN(dz, MAX(0, \frac{\ln(i) - \ln(ik2)}{\eta})) \quad (3.323)$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|--|--------|
| i | Light (PAR) in actual layer | mol photon m ⁻² d ⁻¹ | A |
| lk2 | Light saturation temp. corrected, diatoms | mol photon m ⁻² d ⁻¹ | A |
| eta | Vertical light extinction in layer | m ⁻¹ | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

zk3: Light penetration (availability) in actual layer for cyanobacteria production, m

$$zk1 = MIN(dz, MAX(0, \frac{\ln(i) - \ln(ik3)}{eta})) \tag{3.324}$$

Where:

| Name | Comment | Unit | Type*) |
|------|---|--|--------|
| i | Light (PAR) in actual layer | mol photon m ⁻² d ⁻¹ | A |
| lk3 | Light saturation temp. corrected, cyanobacteria | mol photon m ⁻² d ⁻¹ | A |
| eta | Vertical light extinction in layer | m ⁻¹ | A |
| dz | Height of actual water layer | m | F |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

unh1: potential NH₄ uptake by flagellates, g N m⁻³ d⁻¹

$$unh1 = PC1 * maxupnh * \frac{NH4}{NH4 + hupnh} \quad (3.325)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC1 | Flagellate C | g C m ⁻³ | S |
| maxupnh | Max. N uptake by phytoplankton during N limitation | g N g C ⁻¹ d ⁻¹ | C |
| hupnh | Half-saturation constant for NH ₄ , phytoplankton uptake | g N m ⁻³ | C |
| NH4 | NH ₄ -N in water | g N m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

unh2: potential NH₄ uptake by diatoms, g N m⁻³ d⁻¹

$$unh2 = PC2 * pda2 * maxupnh * \frac{NH4}{NH4 + hupnh * pdb2} \quad (3.326)$$

Where:

$$pda2 = \left(\frac{esd2}{esd1}\right)^{kbt1} \quad (3.327)$$

And:

$$pdb2 = \left(\frac{esd2}{esd1}\right)^{kbt2} \quad (3.328)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC2 | Diatom C | g C m ⁻³ | S |
| maxupnh | Max. N uptake by flagellates during N limitation | g N g C ⁻¹ d ⁻¹ | C |
| hupnh | Half-saturation constant for NH ₄ , phytoplankton uptake | g N m ⁻³ | C |
| NH4 | NH ₄ -N in water | g N m ⁻³ | S |
| pda2 | Ratio, nutrient uptake, Diatom : Flagellate | n.u. | A |
| pdb2 | Ratio, half saturation conc. Diatom: Flagellate | n.u. | A |
| esd1 | Equivalent spherical diameter, flagellates | µm | C |

| Name | Comment | Unit | Type*) |
|-------|--|------|--------|
| esd2 | Equivalent spherical diameter, Diatom | µm | C |
| kbet1 | Exponent 1 for potential uptake of nutrients | n.u. | C |
| kbet2 | Exponent 2 for half saturation conc. | n.u. | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

unh3: Potential NH4 uptake by cyanobacteria, g N m⁻³ d⁻¹

$$unh3 = PC3 * pda3 * maxupnh * \frac{NH4}{NH4 + hupnh * pdb3} \tag{3.329}$$

Where:

$$pda3 = \left(\frac{esd3}{esd1}\right)^{kbt1} \tag{3.330}$$

And:

$$pdb3 = \left(\frac{esd3}{esd1}\right)^{kbt2} \tag{3.331}$$

Where:

| Name | Comment | Unit | Type*) |
|---------|--|---------------------------------------|--------|
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| maxupnh | Max. N uptake by flagellates during N limitation | g N g C ⁻¹ d ⁻¹ | C |
| hupnh | Half-saturation constant for NH4, phytoplankton uptake | g N m ⁻³ | C |
| NH4 | NH4-N in water | g N m ⁻³ | S |
| pda3 | Ratio, nutrient uptake, Cyanobacteria : Flagellate | n.u. | A |
| pdb3 | Ratio, half saturation conc. Cyanobacteria: Flagellate | n.u. | A |
| esd1 | Equivalent spherical diameter, flagellates | µm | C |
| esd3 | Equivalent spherical diameter, Cyanobacteria | µm | C |
| kbet1 | Exponent 1 for potential uptake of nutrients | n.u. | C |
| kbet2 | Exponent 2 for half saturation conc. | n.u. | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

un31: Potential NO₃ uptake by flagellate, g N m⁻³ d⁻¹

$$un31 = PC1 * maxupn3 * \frac{NO3}{NO3 + hupn3} \quad (3.332)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC1 | Flagellate C | g C m ⁻³ | S |
| maxupn3 | Max. NO ₃ uptake by phytoplankton during N limitation | g N g C ⁻¹ d ⁻¹ | C |
| hupn3 | Half-saturation constant for NO ₃ , phytoplankton uptake | g N m ⁻³ | C |
| NO3 | NO ₃ -N in water | g N m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

un32: Potential NO₃ uptake by diatoms, g N m⁻³ d⁻¹

$$un32 = PC2 * pda2 * maxupn3 * \frac{NO3}{NO3 + hupn3 * pdb2} \quad (3.333)$$

Where:

$$pda2 = \left(\frac{esd2}{esd1}\right)^{kbt1} \quad (3.334)$$

And:

$$pdb2 = \left(\frac{esd2}{esd1}\right)^{kbt2} \quad (3.335)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC2 | Diatom C | g C m ⁻³ | S |
| maxupn3 | Max. NO ₃ uptake by flagellates during N limitation | g N g C ⁻¹ d ⁻¹ | C |
| hupn3 | Half-saturation constant for NO ₃ , phytoplankton uptake | g N m ⁻³ | C |
| NO3 | NO ₃ -N in water | g N m ⁻³ | S |
| pda2 | Ratio, nutrient uptake, Diatom : Flagellate | n.u. | A |
| pdb2 | Ratio, half saturation conc. Diatom: Flagellate | n.u. | A |
| esd1 | Equivalent spherical diameter, flagellates | µm | C |
| esd2 | Equivalent spherical diameter, Diatom | µm | C |

| Name | Comment | Unit | Type*) |
|-------|--|------|--------|
| kbet1 | Exponent 1 for potential uptake of nutrients | n.u. | C |
| kbet2 | Exponent 2 for half saturation conc. | n.u. | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

un33: Potential NH₄ uptake by cyanobacteria, g N m⁻³ d⁻¹

$$un33 = PC3 * pda3 * maxupn3 * \frac{NO3}{NO3 + hupn3 * pdb3} \tag{3.336}$$

Where:

And:

$$pda3 = \left(\frac{esd3}{esd1}\right)^{kbt1} \tag{3.337}$$

$$pdb3 = \left(\frac{esd3}{esd1}\right)^{kbt2} \tag{3.338}$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| maxupn3 | Max. NO ₃ uptake by flagellates during N limitation | g N g C ⁻¹ d ⁻¹ | C |
| Hupn3 | Half-saturation constant for NO ₃ , phytoplankton uptake | g N m ⁻³ | C |
| NO3 | NO ₃ -N in water | g N m ⁻³ | S |
| pda3 | Ratio, nutrient uptake, Cyanobacteria : Flagellate | n.u. | A |
| pdb3 | Ratio, half saturation conc. Cyanobacteria: Flagellate | n.u. | A |
| esd1 | Equivalent spherical diameter, flagellates | µm | C |
| esd3 | Equivalent spherical diameter, Cyanobacteria | µm | C |
| kbet1 | Exponent 1 for potential uptake of nutrients | n.u. | C |
| kbet2 | Exponent 2 for half saturation conc. | n.u. | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

upo1: Potential PO₄ uptake by flagellate, g P m⁻³ d⁻¹

$$upo1 = PC1 * maxupip * \frac{PO4}{PO4 + hupp} \quad (3.339)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC1 | Flagellate C | g C m ⁻³ | S |
| maxupip | Max. P uptake by phytoplankton during P limitation | g P g C ⁻¹ d ⁻¹ | C |
| hupip | Half-saturation constant for PO ₄ , phytoplankton uptake | g P m ⁻³ | C |
| PO4 | PO ₄ -P in water | g P m ⁻³ | S |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

upo2: Potential PO₄ uptake by diatoms, g P m⁻³ d⁻¹

$$upo2 = PC2 * pda2 * maxupip * \frac{PO4}{PO4 + hupp * pdb2} \quad (3.340)$$

Where:

$$pda2 = \left(\frac{esd2}{esd1}\right)^{kbt1} \quad (3.341)$$

And:

$$pdb2 = \left(\frac{esd2}{esd1}\right)^{kbt2} \quad (3.342)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC2 | Diatom C | g C m ⁻³ | S |
| maxupip | Max. PO ₄ uptake by flagellates during P limitation | g P g C ⁻¹ d ⁻¹ | C |
| hupp | Half-saturation constant for PO ₄ , phytoplankton uptake | g P m ⁻³ | C |
| PO4 | PO ₄ -P in water | g P m ⁻³ | S |
| pda2 | Ratio, nutrient uptake, Diatom : Flagellate | n.u. | A |
| pdb2 | Ratio, half saturation conc. Diatom: Flagellate | n.u. | A |
| esd1 | Equivalent spherical diameter, flagellates | µm | C |

| Name | Comment | Unit | Type*) |
|-------|--|------|--------|
| esd2 | Equivalent spherical diameter, Diatom | µm | C |
| kbet1 | Exponent 1 for potential uptake of nutrients | n.u. | C |
| kbet2 | Exponent 2 for half saturation conc. | n.u. | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

upo3: Potential PO₄ uptake by cyanobacteria, g P m⁻³ d⁻¹

$$upo3 = PC3 * pda3 * maxupip * \frac{PO4}{PO4 + hupp * pdb3} \quad (3.343)$$

Where:

$$pda3 = \left(\frac{esd3}{esd1}\right)^{kbt1} \quad (3.344)$$

And:

$$pdb3 = \left(\frac{esd3}{esd1}\right)^{kbt2} \quad (3.345)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|---------------------------------------|--------|
| PC3 | Cyanobacteria C | g C m ⁻³ | S |
| maxupip | Max. PO ₄ uptake by flagellates during P limitation | g P g C ⁻¹ d ⁻¹ | C |
| hupp | Half-saturation constant for PO ₄ , phytoplankton uptake | g P m ⁻³ | C |
| PO4 | PO ₄ -P in water | g P m ⁻³ | S |
| pda3 | Ratio, nutrient uptake, Cyanobacteria : Flagellate | n.u. | A |
| pdb3 | Ratio, half saturation conc. Cyanobacteria: Flagellate | n.u. | A |
| esd1 | Equivalent spherical diameter, flagellates | µm | C |
| esd3 | Equivalent spherical diameter, Cyanobacteria | µm | C |
| kbet1 | Exponent 1 for potential uptake of nutrients | n.u. | C |
| kbet2 | Exponent 2 for half saturation conc. | n.u. | C |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

usi2: Potential Si uptake by diatoms, g Si m⁻³ d⁻¹

$$usi2 = PC2 * maxupsi * \frac{Six}{Six + hupsi} \quad (3.346)$$

Where:

$$six = Si - Simin \quad (3.347)$$

Where:

| Name | Comment | Unit | Type*) |
|---------|---|--|--------|
| PC2 | Flagellate C | g C m ⁻³ | S |
| maxupsi | Max. Si uptake by phytoplankton during Si limitation | g Si g C ⁻¹ d ⁻¹ | C |
| hupsi | Half-saturation constant for Si, diatom uptake | g Si m ⁻³ | C |
| Six | available Si for diatoms, (Si-Simin) >=0, Si for uptake PC2 | g Si m ⁻³ | A |
| Si | Si in water | g Si m ⁻³ | S |
| Simin | Si not available for PC2 | g Si m ⁻³ | A |

*) S: State variable, F: Forcing, C: Constant, P: Process in differential equation, P1: Help process, A: Auxiliary help process.

4 Data Requirements

- Basic Model Parameters
 - Model grid size and extent
 - Time step and length of simulation
 - Type of output required and its frequency
- Bathymetry and Hydrodynamic Input
- Combined Advection-Dispersion Model
 - Dispersion coefficients
- Initial Conditions
 - Concentration of parameters
- Boundary Conditions
 - Concentration of parameters
- Pollution Sources
 - Discharge magnitudes and concentration of parameters
- Process Rates
 - Size of coefficients governing the process rates. Some of these coefficients can be determined by calibration. Others will be based on literature values or found from actual measurements and laboratory tests.
- Forcings
- Data sets of photosynthetic active light (PAR) ($E/m^2/day$)

5 References

- /1/ Droop M.R. 1973. Some thoughts on nutrient limitation in algae. *J. Phycol.* 9 P:264-272.
- /2/ Droop M.R. 1975. The nutrient status of algae cells in batch cultures. *J. Mar. Biol. Ass. U.K.* 55 P:541-555.
- /3/ Effer S.W. 1988. Secchi disk transparency and turbidity. *Journal of environmental engineering.* 1988 vol. 114 no. 6 pp. 1436-1447.
- /4/ Kirk J.T.O. 2000. *Light and photosynthesis in aquatic ecosystems.* Cambridge University Press 2. Edition 2000. ISBN 0521453534, ISBN 052145459664
- /5/ Lund-Hansen L.C. 2004. Diffuse attenuation coefficients $K_d(\text{PAR})$ at estuarine North Sea-Baltic Sea transition: time-series, partitioning, absorption, and scattering. *Estuarine, Coastal and Shelf Science* 61 (2004) pp:251-259.
- /6/ MIKE by DHI 2011a. MIKE 21 & MIKE 3 Flow Model. Mud Transport module Scientific Description. DHI water environment health, Hørsholm Denmark
- /7/ MIKE by DHI 2011b. ECO Lab User guide. DHI water environment health, Hørsholm Denmark
- /8/ Monod J. 1949. The Growth of Bacterial Cultures. *Annual Review of Microbiology*, v. 3, p. 371.
- /9/ Morel A. 1980. In water and remote measurements of ocean color. *Boundary-Layer Meteorology* 18 (1980) pp 177-201.
- /10/ Nyholm N. 1977 Kinetics of phosphate-limited algae growth. *Biotechn. Bioengineering* 19 P:467-492.
- /11/ Nyholm N. 1978 A simulation model for phytoplankton growth cycling in eutrophic shallow lakes. *Ecological Modelling.* Vol. 4, P:279-310.
- /12/ Nyholm N. 1979 The use of management models for lakes at the Water Quality Institute. Denmark. *State-of-the-art in Ecological Modelling.* Vol. 7. P:561-577.
- /13/ Pfannkuche F. 2002. Optical properties of Otago Shelf Waters: South Island New Zealand. *Estuarine, Coastal and Shelf Science* 55 (2002) pp:613-627.
- /14/ Prieur L., S. Sathyendranath 1981. An optical classification of coastal and oceanic waters based on the specific spectral absorption curves of phytoplankton pigments, dissolved organic matter, and other particulate materials. *Limnol. Oceanogr.* 26(4) PP 671-689.
- /15/ Rasmussen Erik Kock, O.S. Petersen, J.R. Thomsen, R. J. Flower, F. Aysche, M. Kraiem, L. Chouba 2009. Model analysis of the future water quality of the eutrophicated Ghar EL Melh lagoon (Northern Tunisia). *Hydrobiologia* (2009) 622:173-193
- /16/ Tett P., A. Edvards & K. Jones 1986. A model for the growth of shelf-sea phytoplankton in summer. *Estuar. Coast. Shelf Sci.* 23 P:641-672.

